

# Journal of Materials Chemistry B

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: J. J. Virgen-Ortíz, J. C.S DOS SANTOS, Á. Berenguer-Murcia, O. Barbosa, R. C. RODRIGUES and R. Fernandez-Lafuente, *J. Mater. Chem. B*, 2017, DOI: 10.1039/C7TB01639E.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [author guidelines](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the ethical guidelines, outlined in our [author and reviewer resource centre](#), still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

# Polyethylenimine: A very useful ionic polymer in the design of immobilized enzyme biocatalysts

Jose J. Virgen-Ortíz<sup>1</sup>, José C.S. dos Santos<sup>2</sup>, Ángel Berenguer-Murcia<sup>3</sup>, Oveimar Barbosa<sup>4</sup>, Rafael C. Rodrigues<sup>5</sup>, Roberto Fernandez-Lafuente<sup>6,\*</sup>

<sup>1</sup>CONACYT - Centro de Investigación en Alimentación y Desarrollo, A.C. (CIAD) - Consorcio CIDAM, Km. 8 Antigua Carretera a Pátzcuaro s/n, 58341 Morelia, Michoacán, México.

<sup>2</sup>Instituto de Engenharias e Desenvolvimento Sustentável, Universidade da Integração Internacional da Lusofonia Afro-Brasileira, CEP 62785-000, Acarape, CE, Brazil.

<sup>3</sup>Instituto Universitario de Materiales, Departamento de Química Inorgánica, Universidad de Alicante, Campus de San Vicente del Raspeig, Ap. 99 - 03080 Alicante, Spain.

<sup>4</sup>Departamento de Química, Facultad de Ciencias. Universidad del Tolima, Ibagué, Colombia.

<sup>5</sup>Biocatalysis and Enzyme Technology Lab, Institute of Food Science and Technology, Federal University of Rio Grande do Sul, Av. Bento Gonçalves, 9500, P.O. Box 15090, Porto Alegre, RS, Brazil.

<sup>6</sup>ICP-CSIC. Campus UAM-CSIC. Cantoblanco. 28049 Madrid. Spain.

\* Corresponding author: Prof. Dr. Roberto Fernández-Lafuente. Departamento de Biocatálisis. ICP-CSIC. C/ Marie Curie 2. Campus UAM-CSIC. Cantoblanco. 28049 Madrid, Spain. e-mail: [rfl@icp.csic.es](mailto:rfl@icp.csic.es).

## Abstract

This review discusses the possible roles of polyethylenimine (PEI) in the design of improved immobilized biocatalysts from diverse perspectives. This includes their use to activate supports and immobilize enzymes via ion exchange, as well as to improve immobilized enzymes by coating with PEI. PEI is a polymer containing primary, secondary and tertiary amino groups, having a strong anion exchange capacity under a broad range of conditions, and the capability to chemically react with different moieties on either enzyme or support. Also, as a multifunctional polymer, it has been modified stepwise to introduce different functionalities in the same polymer. This polymer (in combination with other anionic ones) permits the generation of “saline” environments around enzyme molecules, improving enzyme stability in the presence of hydrophobic compounds. The use of PEI as a physical glue useful to crosslink enzymes subunits in multimeric enzymes, monomeric enzymes immobilized via physical interactions or production of enzyme multilayers will be specially emphasized as new open avenues for enzyme coimmobilization. The coimmobilization of enzymes and cofactors using PEI may become one of the future developments allowed through an adequate use of this polymer and new pathways towards the design of enzyme combi-catalysts for their use in cascade reactions. Some unexplored but suggested uses derived from the properties of PEI are also proposed in the review, like the use of the buffering power of this multifunctional polymer to avoid pH gradients inside the biocatalyst particle. Thus, although PEI has been a largely popular polymer in biocatalysts design, it looks that a long and in some cases almost unexplored road lies ahead.

**Key words:** generation of nanoenvironments, physical crosslinking, multimeric enzymes, coimmobilization of enzymes and cofactors, CLEAs crosslinking, cascade reactions

## 1. Introduction: The role of immobilization and physicochemical modification of enzymes in the design of industrial biocatalysts

One of the main goals of chemistry in the 21<sup>st</sup> century is the design of greener and more environmentally sustainable processes.<sup>1-3</sup> Enzyme biocatalysis has risen as a very suitable pathway towards this goal;<sup>4</sup> enzyme catalysis is selective, specific and may be performed under mild conditions.<sup>5-8</sup> However, enzymes have evolved to fulfill the natural requirements of microorganisms, e.g. capacity to respond to changes in the medium or adaptation to stress conditions, and they have many properties that greatly differ from those required from an industrial catalyst.<sup>9-11</sup> For example, enzymes are moderately stable, and may be inhibited by products, substrates or even non-related compounds.<sup>12</sup> One additional problem is that most proteins are water soluble, making recovery of this expensive catalyst complex (almost impossible in some cases). These problems have hampered their industrial implementation. Nowadays, revolutions in many areas related to the design of biocatalysts, such as metagenomics,<sup>13,14</sup> directed evolution,<sup>15,16</sup> or site-directed mutagenesis,<sup>17</sup> have reduced the price of enzymes and have thus increased their availability. In this review, we will focus on enzyme immobilization. Enzyme immobilization needs to be performed to facilitate enzyme recovery (Figure 1). This prevents product contamination and favors enzyme reuse, provided that the immobilized enzyme is stable enough to be reused.<sup>18-20</sup>

The necessity for enzyme immobilization has spurred an intensive research to couple immobilization to the improvement of other enzyme features. It has been shown that a proper immobilization may improve enzyme stability (e.g., via multipoint or multisubunit immobilization (Figures 2 and 3), generation of suitable enzyme nano-environments (Figure 4)), but may also improve enzyme activity (by generating more active conformations or using internal substrates or products

69 gradients), specificity or selectivity.<sup>9,21–26</sup> Even enzyme purity may be enhanced using a properly  
70 designed immobilization protocol, saving time, avoiding side reactions that can decrease process yields  
71 or selectivity/specificity, and decreasing the amount of support required to immobilize a fixed amount  
72 of the target enzyme.<sup>27</sup> All these enzyme features are related in a certain sense. For example, as stated  
73 by J. Woodley,<sup>28</sup> from a practical point of view, enzyme stability should be considered not just as an  
74 operational lifetime indicator, but should be measured as the amount of synthesized product per  
75 biocatalyst mass (price of product/price of used biocatalyst). This means that if immobilization of the  
76 enzyme increases the operational half-life, and also reduces the reaction time course (e.g. due to a  
77 reduction of inhibition problems or an improved enzyme activity), the “stability” of the enzyme will be  
78 boosted by all these improvements.

79 However, enzyme immobilization has some costs that must be compensated for by the gains:  
80 support and processing costs. While much attention has been paid to the support costs (even though it  
81 is much cheaper than the enzyme), it should be considered that if a support that has a higher cost (let us  
82 suppose by a 10-fold factor) permits a higher loading, a higher enzyme stability and productivity, its  
83 price may become irrelevant. Moreover, reversible enzyme immobilization methods have the  
84 advantage of reducing the weight of the support price in the immobilized enzyme biocatalysts, because  
85 the support may be reused after enzyme inactivation: the inactivated enzyme may be released and fresh  
86 enzyme may be added (Figure 1).<sup>23</sup> The main point is to have so strong an enzyme reversible  
87 immobilization that the enzyme cannot become desorbed during operation. Physical adsorption is the  
88 simplest reversible immobilization,<sup>29</sup> but it is not devoid problems because the support surface will  
89 always be physically active and that can generate some problems regarding enzyme support  
90 uncontrolled interactions.<sup>30</sup> Furthermore it is not possible to design approaches for protein reactivation  
91 using unfolding/refolding strategies.<sup>23</sup> However, in certain cases, physical adsorption of the enzyme  
92 may produce some enzyme stabilization, like in the case of multimeric enzymes, if this immobilization

involves all enzyme subunits and prevents enzyme dissociation (Figures 2 and 3).<sup>24</sup> Interfacial activation of lipases on hydrophobic supports<sup>31</sup> is perhaps one of the most successful examples where physical adsorption may stabilize biocatalysts even to a larger degree than multipoint covalent attachment<sup>32,33</sup> due to the existence of a specific enzyme form in these supports that is more stable than the lipase in equilibrium conformation.<sup>34</sup> However, even in this case, the enzyme may be desorbed during operation and reduce the application of this method.<sup>35,36</sup>

Physical or chemical modification of enzymes is another way to improve enzyme properties (Figure 5).<sup>37,38</sup> The introduction of intramolecular crosslinkings with small bifunctional reagents may greatly increase enzyme rigidity,<sup>39,40</sup> while inter-subunit crosslinking of multimeric enzymes may avoid subunit dissociation (Figures 2 and 3).<sup>24</sup> There are many examples where the enzyme properties (activity, stability, specificity) have been improved by a global modification of the enzyme surface (Figure 5).<sup>41–44</sup> Immobilization and physicochemical modification of enzymes are not unrelated tools to improve enzyme properties, but may be used in a coupled way to improve the final performance of the enzyme biocatalyst (Figure 5).<sup>45,46</sup> In many instances, immobilization of the enzyme may facilitate further modifications (chemical or physical). In other cases, chemical modification is designed to improve enzyme immobilization (Figure 3).<sup>47</sup>

In this review, the use of polyethylenimine as a reagent for the preparation of immobilized enzyme biocatalysts will be reviewed, both as a method to activate supports as ion exchangers or to physically modify enzymes before or after their immobilization. This multifunctional polymer has been employed in biocatalysis design in many instances and with very different objectives. The properties of this polymer and some of their applications will be presented and discussed, e.g., stabilization of multimeric enzymes (Figures 2 and 3), stabilization of enzymes versus organic solvents or oxygen (Figure 6), immobilization of enzyme multilayers (e.g., coimmobilization of enzymes) (Figure 7) or coimmobilization of enzymes and cofactors. The potential of this polymer in biocatalyst

design is already quite impressive and considering the latest applications, it is safe to assume that it has not been fully exploited yet.

## 2. Production and features of PEI

Polyethylenimine (PEI) is one of the most widely used synthetic polycations in various applications because of its chemical functionality arising from the presence of cationic primary (25%), secondary (50%), and tertiary amines (25%).<sup>48,49</sup> PEI is formed by the linking of iminoethylene units and can have linear, branched, comb, network, and dendrimer architectures depending upon its synthesis and modification methods, which greatly influences its properties, both physical and chemical.<sup>50</sup> Furthermore, these synthetic approaches enable PEI to be available in a wide range of molecular weights. At room temperature, branched PEI (BPEI) is a highly viscous liquid while linear PEI (LPEI) is a solid. PEI has several attractive features for its use in widespread applications, such as low toxicity, ease in its separation and recycling, and (last but not least) its being odorless. In addition to these attractive features, there is a distinct feature of PEI which places it ahead of other polyions (e.g. polyallilamine or chitosan) when it comes to loading, and which justifies its widespread use in fields as varied as detergents, adhesives, water treatment, cosmetics, carbon dioxide capture,<sup>51–54</sup> as DNA transfection agent, and in drug delivery:<sup>55–58</sup> despite being a weak polymeric base with  $pK_a$  values between 7.9 and 9.6, it possesses a high ionic charge density, which in practical terms translates in being a more cost-effective material. This derives in either the possibility of reaching the same loadings with reduced amounts of the polymer (which would colloquially mean “getting a bigger bang for the buck”) or reaching loadings which are beyond the reach of the aforementioned examples while avoiding enzyme agglomeration thanks to its multibranched network.

Commercial polyethylenimine usually has a branched structure. BPEI is soluble in polar solvents as water, ethanol, and acetone and depending on the molecular weight is partially soluble in benzene and tetrahydrofuran<sup>59,60</sup>. This molecule will be the most interesting in biocatalysts design, with a random structure and a capability of covering the surface of supports or proteins. The standard synthetic route for obtaining BPEI involves the electrophilic (or cationic) ring-opening polymerization of unsubstituted ethylenimine with a protonic catalyst such as Lewis acids and their salts, strong Brønsted acids, alkyl halides, strong acid esters, phenols, carboxylic acids, or halogens (Scheme 1).<sup>60</sup> This polymerization causes the formation of a secondary amino group in the polymer chain which is more basic than the monomer. Therefore, chain transfer to the polymer takes place extensively, which ultimately brings about branching. The branched structure from ethylenimine polymerization causes the spheroidal shaped of the BPEI molecules.

Another route for obtaining BPEI contemplates the ring-opening isomerization polymerization of substituted 2-oxazolines with Lewis acids, protonic acids, sulfonate esters, or sulfonic anhydrides as inductors the polymerization (Scheme 2). The reaction rate and yield depend on the substituent, the nature of X<sup>-</sup>, solvent, and temperature.<sup>49,60</sup>

On the other hand, linear polyethylenimine is highly crystalline due to its linear structure. Through X-ray studies it was determined that LPEI forms double-stranded helical chains stabilized by interchain hydrogen bonds. The polymer is remarkably hygroscopic, which is due to a transformation from the double stranded helix to a fully extended form upon water adsorption of water. The production of LPEI does not proceed through the ring-opening route of ethylenimine (aziridine) because very high reaction times are required and the yields of the obtained product are very low. Thence, the preparation of LPEI is carried out via ring-opening polymerization (ROP) of 2-ethyl-2-oxazoline followed by hydrolysis, where it is possible to obtain 100% conversion under appropriate conditions (Scheme 3).<sup>48,60</sup>



### 3. Use of PEI to immobilize enzymes and proteins on pre-existing supports

Activation of supports by coating their surface with PEI enables enzyme immobilization by anion exchange. This specific use of PEI was reviewed in 1991,<sup>61</sup> but many advances have been made since then. This immobilization protocol requires the possibility of forming several ion bonds between the enzyme and the support, shifting the counter ions that both surfaces will have.<sup>62</sup>

Immobilization of biomacromolecules using these supports having polymeric beds presents some advantages compared to standard anion exchangers.<sup>61,63</sup> Firstly, enzyme adsorption is quite strong, as PEI has multiple cation groups at different distances that may be adapted to the distance between the enzyme groups. Secondly, the polymer is random coil and will not force the enzyme to become distorted when interacting with it via multiple points. Third, the polymeric bed formed by PEI offers a three-dimensional adsorption versus the two-dimensional adsorption of standard supports,<sup>64</sup> enabling the immobilization of more than 80% of the proteins contained in a crude protein extract.<sup>65</sup> However, immobilization remains reversible and the support may be reused (Figure 1).

Considering that supports coated with anionic polymers may also immobilize over 80 % of the proteins under the same conditions,<sup>66</sup> it seems evident that the net charge of the protein is not the reason for enzyme immobilization, but the possibility of establishing many ionic bridges. This idea has permitted the development of different tailor-made ion exchangers for selective immobilization of different proteins.<sup>67</sup>

However, the use of PEI may also raise some problems, e.g., the polymer can interact with internal pockets of the protein (e.g., where some critical cations may be located) leading to enzyme inactivation: this does not occur using small cationic groups on a flat surface. Moreover, the mobility of PEI may give a certain destabilization of the enzyme that is interacting with it as a result.<sup>63</sup> Ion

exchange should not provide any significant structure rigidification of the enzyme, and even less using PEI due to its high mobility, although some stabilizing effects may be found which are not based in enzyme rigidification, as described in other points of this review.

To maximize the reactive bed volume, the way the coating is produced is critical. Conditions that favor the attachment of many PEI molecules will be preferred versus conditions that favor the spreading of the PEI on the support surface (Figure 8).<sup>65</sup> It is possible to allow the enzyme molecule to penetrate deeper into the polymeric bed, using conditions where the adsorption is not favored and where a higher number of enzyme-support ion bridges is required to fix the enzyme to the support,<sup>68</sup> increasing the adsorption strength and yielding other beneficial effects of immobilization in this kind of supports.<sup>61</sup>

This PEI coating has been used in many different supports and applications. Now, we will review some of the main examples.

### **3.1. Use of PEI to coat pre-existing macroporous supports to immobilize proteins**

#### **3.1.1. Use of supports just activated PEI**

Different supports have been activated for a long time with PEI. Using this strategy, it has been found that PEI can have some positive effects on enzyme stability by generating a partition of hydrophobic compounds (like organic solvents or gases like oxygen) from the enzyme environment (Figures 4 and 6).<sup>69</sup> In general, support preparation is not optimized to maximize the polymeric bed thickness, as just one molecule of the polymer may be enough to immobilize a protein, although this may reduce the impact of the immobilization on the enzyme properties. The fact that this support may rapidly adsorb most proteins in crude extract makes the presence of contaminants detrimental since it may reduce the final loading of the target protein. This occurred when using an excess of total

protein,<sup>70</sup> and gave an unexpected evolution of the amount of immobilized protein when increasing the amount of offered enzyme.<sup>22</sup> These activated supports have been used in many instances in the preparation of biocatalysts or biosensors. For example, they have even been utilized to immobilize antibodies in biosensors. This application may give freedom of movement to the antibody and avoid steric hindrances for the recognition of the antigen.<sup>71–77</sup> However, considering the huge potential of PEI-coated supports as anion exchangers, it does not look very attractive, as in these cases an inert surface is recommended to prevent false positive.<sup>78–80</sup>

There are many examples in the literature on the use of supports activated only with PEI to immobilize enzymes as biocatalysts.<sup>81–116</sup> Table 1 offers the main reports.

The most important advantage is the simplicity of the immobilization protocol; the process consists in simply mixing the enzyme and the support (usually at pH 7), giving very high immobilization rate, high immobilization yields and expressed activities. However, stabilization of the enzymes is relatively poor compared to the stabilization that can be achieved using covalent immobilization using short spacer arms. An exception is the immobilization of multimeric enzymes (Figure 2 and 3) [24] or stability in the presence of organic solvents (Figure 6). This is recommended mainly for very stable enzymes or when the support is so expensive that it is convenient to reuse it.

The reinforcement of the adsorption may prevent undesired enzyme desorption during operation. To this goal, genetic<sup>117</sup> or chemical<sup>118</sup> enrichment of the enzyme surface in anionic groups has been utilized to strengthen the ion exchange of penicillin G acylase on supports coated with PEI.

This support may be also utilized to co-immobilize several enzymes to catalyze a cascade reaction. That is the case of the coimmobilization of cyclohexanone monooxygenase and glucose-6-phosphate dehydrogenase to transform (2-oxyocyclohexyl)acetic acid into (2-oxooexpan-2-yl)acetic acid.<sup>119</sup>

In some cases the lay-out of the reactor is very complex and may be a problem if it needs to be dismantled and remounted after enzyme inactivation. In these cases, the reversibility of the immobilization of the enzymes on PEI coated supports become a clear advantage (Figure 9).

One additional problem is that the inactivation of proteins near to the PEI may drive to a fully composites enzyme-polymer where the enzyme maximize the number of enzyme-support interactions(Figure 10)<sup>120,121</sup>. That way, the full desorption of inactivated enzymes immobilized on PEI activated supports may become much harder than the desorption of native proteins. Optimization of this point needs to be considered if the reversibility of the immobilization is a key point to select this protocol.

As stated in the introduction, this ion exchange in a polymeric bed may permit to stabilize multimeric enzymes by involving all enzyme subunits or stabilizing any protein by generating a favorable environment (Figures 2 and 3). However, real rigidification of the enzyme should not be expected (the bonds are weak and the polymer flexible) and the support remains capable of interacting with the protein, which may even produce some enzyme destabilization.<sup>120,121</sup>

### 3.1.2. Chemical crosslinking of enzyme adsorbed on PEI coated supports

In some cases, to further improve the enzyme properties and fully prevent enzyme desorption, the adsorbed enzymes have been treated with bifunctional chemical reagents, usually glutaraldehyde, to get a covalent immobilization. This immobilization method is very versatile, and some considerations should be brought forward in their use, as the possibility of activating each primary amino group with one or two glutaraldehyde molecules, since this may produce fully different effects.<sup>122–125</sup> This strategy means that reversibility is lost, therefore it must be evaluated whether the gains justify this shortcoming or not. The treatment with glutaraldehyde of the adsorbed enzymes will also crosslink the polymer<sup>124</sup> which will become more rigid and perhaps able to transmit some rigidity

to the enzyme. Moreover, the enzyme will be modified by glutaraldehyde, in some instances improving the enzyme stability or activity,<sup>44,123,126–128</sup> but in some others it may have highly detrimental effects on enzyme properties. Table 2 shows some examples of the use of glutaraldehyde to prevent enzyme desorption of enzymes immobilized in PEI coated supports<sup>129–143</sup>.

Other bifunctional reagents have also been utilized to get this covalent crosslinking, like dimethyl suberimidate (in the immobilization of Jack bean urease).<sup>144</sup> Uridine phosphorylase was immobilized on PEI activated support and further treated with aldehyde dextran to stabilize their multimeric structure, greatly improving its stability.<sup>145</sup> A similar strategy was used later to stabilize the multimeric nucleoside 2'-deoxyribosyltransferase from *Bacillus psychrosaccharolyticus*.<sup>146,147</sup>

In a more sophisticated strategy, polymerization of methyl methacrylate, bromoethyl methacrylate and ethylene glycol dimethacrylate was used to produce a bromide-functionalized matrix.<sup>148</sup> The obtained solid particles were utilized as initiators for the ring-opening polymerization of 2-methyl 2-oxazoline. Finally, the surface brushes were converted into polyethylenimine by acid hydrolysis, activated with glutaraldehyde and used to immobilize  $\alpha$ -amylase.<sup>148</sup> The protocol seems to be too complex for industrial implementation, as results did not surpass conventional covalent immobilization. In our view, to use glutaraldehyde or any other crosslinking reagent, will greatly decrease the advantages of PEI, although a higher percentage of the enzyme surface may be involved in the covalent attachment, each bond will have a minimal effect on the enzyme stability due to the polymer flexibility.

### 3.1.3. Use of PEI supports treated with bifunctional reagents to get a covalent enzyme immobilization

In some other cases, the support activated with PEI is treated with glutaraldehyde or other reagents before immobilization of the enzyme to directly obtain the covalent immobilization of the enzyme. Considering that this must produce the inter and intramolecular crosslinking of PEI, the advantages of this protocol compared to the use of really rigid flat surfaces activated with similar levels of activation of small amino groups is unclear; perhaps it is not easy to reach similar levels of activation using standard supports compared to the values achieved using PEI. Table 3 shows some of the examples using glutaraldehyde as support activating reagent<sup>149-149-171</sup>.

Other bifunctional reagents have been used to activate PEI, like dimethyl sulphate or triethyloxonium tetrafluoroborate to immobilize pectinesterase,<sup>172</sup> epichlorohydrin to immobilize catalase<sup>173</sup> and cholesterol oxidase.<sup>174</sup> Poly-functional aldehyde-dextran has been used to get this covalent crosslinking, like in the case of the nucleoside 2'-deoxyribosyltransferase from *Bacillus psychrosaccharolyticus*.<sup>146</sup> In some cases, unfortunately quite scarce, a comparison among different PEI activation methods is presented. For example, one paper reports that PEI (in some instances partially modified with succinic anhydride) was activated with glutaraldehyde or hexamethylene diisocyanate to immobilize lipase from *T. lanuginosus*.<sup>175</sup> A higher stability was achieved using glutaraldehyde.

In other cases more sophisticated strategies were employed to get covalent attachments. For example, PEI activated supports were treated with epichlorohydrin and then modified with thiophosgene or with succinic anhydride; the carboxyl groups were then converted to hydrazide. These supports were successfully used to immobilize D-glucose oxidase, glucoamylase and cholinesterase (using hydrazide supports the sugar chains of the enzymes were submitted to periodate oxidation).<sup>176</sup> In another case, PEI was modified with aryldiazirine derivatives. After UV irradiation the groups were active (diazirine) and were used to immobilize lysozyme, pepsin, and horseradish peroxidase on

polypropylene films.<sup>177</sup> The sophistication of these methods makes them complex to implement at industrial level, although they have opened new opportunities should the need arise.

The flexibility of PEI was a key point in the design of different biosensors where the transmission of the electron is a requirement for its proper performance. For example, glucose oxidase and ferrocene were immobilized on PEI to detect glucose.<sup>178</sup> The multifunctionality of PEI is one of the advantages of using this interesting polymer in biocatalysts design, as stepwise modifications are possible.

### 3.2. Use of PEI to activate nanoparticles to immobilize enzymes

#### 3.2.1. Use of nanoparticles coated with PEI to immobilize enzymes

Immobilization of enzymes in nanoparticles is growing in interest in the scientific literature (Figure 11).<sup>179–181</sup> Non-porous nanoparticles have some advantages, mainly related to the decrease of the diffusional problems as all the enzyme will be on their surface.<sup>23</sup> This makes that this immobilization method may be recommended for using immobilized enzymes versus solid or very large substrates, or even just in the presence of a suspension although the substrate itself may be soluble (e.g., in fruit juice) (Figure 11).<sup>23</sup> In cases where the activity of the enzyme is very high, nonporous nanoparticles prevent the existence of gradients (pH gradients, substrate gradients, even inactivating reagents like H<sub>2</sub>O<sub>2</sub> gradients) that can affect the enzyme features,<sup>182–184</sup> not always in a negative way.<sup>22,185</sup> Using porous supports coated with PEI, the cationic polymer could behave like a solid buffer, and could prevent pH gradients formation, but we have not found any report regarding this fact (Figure 12).

However, immobilization in nonporous supports also has disadvantages: the enzyme is exposed to the medium and may be inactivated by proteolysis (catalyzed by protease molecules immobilized on other particles), or it may interact with hydrophobic interfaces (e.g., gas bubbles) (Figure 11), etc.<sup>23</sup>

Coating with polymers has been proposed to solve this problem, and proteins coated with either aldehyde dextran<sup>186,187</sup> or PEI<sup>188,189</sup> have been proved to be more resistant to this kind of inactivation. Thus, enzymes immobilized inside a polymeric PEI bead could benefit from this protective effect.

The handling of nanomaterials may be complex. This has been solved using para-magnetic materials. Magnetic nanoparticles may be easily recovered from the reaction medium using a magnet and that solves the handling problem of nanoparticles use.<sup>190,191</sup> Moreover, nanomaterials tend to be expensive, and the use of reversible immobilization methods could permit reusing the support. PEI-coated magnetic nanoparticles have been claimed as adequate for enzyme immobilization and present limited environmental impact.<sup>192</sup> Thus, there are many examples of immobilization of proteins in nanoparticles using supports coated with PEI (Figure 11).

The first examples found on this kind of immobilization dated from 2007, as the development of nanotechnology was a requirement prior to this application.<sup>193</sup> Lipases are among the enzymes most usually immobilized on nanoparticles activated with PEI.<sup>194</sup> Superparamagnetic Fe<sub>3</sub>O<sub>4</sub> particles coated with PEI were used to immobilize and partially purify lipase from *Candida rugosa*.<sup>195,196</sup> Later, this enzyme was immobilized on zinc oxide (ZnO) nanoparticles with PEI activated with glutaraldehyde and utilized in the synthesis of geranyl acetate.<sup>197</sup> Lipase from *T. lanuginosus* was immobilized on polyethyleneimine-modified superparamagnetic Fe<sub>3</sub>O<sub>4</sub> nanoparticles and further crosslinked using glutaraldehyde.<sup>198</sup> This enzyme was later immobilized in a similar support assisted with divalent metal chelated ions.<sup>199</sup> Lipases have been immobilized on PEI-modified magnetic nanoparticles and used to produce vitamin A palmitate via esterification reactions.<sup>200</sup> Fe<sub>3</sub>O<sub>4</sub> magnetic nanoparticles were coated with PEI followed by grafting  $\beta$ -cyclodextrin and utilized to immobilize lipase utilized in the production of ethyl valerate.<sup>201</sup> Other enzymes have also been immobilized on PEI coated nanoparticles.  $\beta$ -galactosidase from *Kluyveromyces fragilis* has been immobilized on magnetic poly(glycidyl methacrylate-ethylene glycol dimethacrylate-hydroxyethyl methacrylate) nanospheres



grafted with polyethyleneimine.<sup>202</sup>  $\alpha$ -rhamnosyl- $\beta$ -glucosidase from *Acremonium sp.* DSM 24697 was immobilized by ion exchange and cross-linking onto PEI-iron particles, improving the results reported using polyanilide.<sup>203</sup> Partially phosphonated polyethylenimine has been used to coat iron oxide nanoparticles and used to covalently immobilize trypsin and used in proteomic analysis.<sup>204</sup> In another paper, metal ion-chelated linear PEIs coating nanoparticles were used to immobilize and stabilize glycerol dehydrogenase.<sup>205</sup> Laccase has also been immobilized on magnetic Fe<sub>3</sub>O<sub>4</sub> nanoparticles modified with polyethylenimine.<sup>206</sup> Cellulose nanocrystals were successfully combined with PEI modified Fe<sub>3</sub>O<sub>4</sub> nanoparticles and successfully used for the immobilization of papain.<sup>207</sup> Acetylcholinesterase has been immobilized on PEI-coated silica nanoparticles improving its stability.<sup>208</sup>

### 3.2.2. Use of PEI to form the nanoparticles to be used as carries of enzymes

In some instances, PEI is not only used to immobilize the enzyme, but also plays a fundamental role in nanoparticle building. PEI is in many cases used as template and reducing agent in the production of nanoparticles. Thus, PEI has been used as template for the formation of silica nanoparticles trapping chemically modified anionic peroxidase, showing direct electron transfer at 0 mV versus Ag/AgCl.<sup>209</sup> Amine dehydrogenase was immobilized on hybrid titania nanoparticles, which were produced by polyethylenimine coating and templated biomineralization.<sup>210</sup>

Other examples where the role of PEI in the nanoparticle may go beyond immobilization of enzymes is the formation of colloidal stable nanoparticles using mixtures of PEI and anionic polymers. For example, mixtures of PEI and poly(maleic acid-co-propylene) have been used to form stable colloidal nanoparticles, crosslinked via maleic anhydride and utilized to immobilized laccase.<sup>211</sup>

Some sophisticated strategies have been used, based on PEI properties to bind the nanoparticles to other solids. For example, submicron-sized poly(N-isopropyl acrylamide)/polyethyleneimine core-

shell microgels were prepared in aqueous media using tert-butyl hydroperoxide as an initiator followed by the formation of nanoparticles on the surface of the microgels.<sup>193</sup> PEI was used as binder for the assembly of the gold nanoparticles/microgel complex, and finally horseradish peroxidase and urease were successfully immobilized.<sup>193</sup> In another paper, bacteria cellulose nanofiber nanocomposites were produced and coated with gold nanoparticles using PEI as reducing and linking agent.<sup>212</sup> Then, this composite was used to immobilize horseradish peroxidase. A similar protocol was utilized later to prepare a support where glucose oxidase was immobilized using carbodiimide as enzyme linking agent in this case.<sup>213</sup> Gold nanoparticles were also immobilized on the inner wall at the inlet end of a capillary electrophoresis unit treated with PEI; and then used to immobilize L-glutamic dehydrogenase.<sup>214</sup> This immobilized enzyme was used as a sensor for enzyme inhibitors. The immobilization of multi-walled carbon nanotubes using nafion and polyethylenimine as dispersants and glucose oxidase as enzyme was utilized to improve the determination of glucose.<sup>215</sup> Self-polymerization of dopamine on the surface of CaCO<sub>3</sub> particles was used to form polydopamine and used as an adhesive layer to bind PEI through Michael addition reaction or Schiff base reaction. Then, titania was generated and deposited on the surface of the PEI nanoparticles via bioinspired mineralization process initiated by the free amine groups of PEI. EDTA treatment produced hybrid microcapsules by eliminating the CaCO<sub>3</sub> templates and used to immobilize catalase.<sup>216</sup> Gold nanoparticles were *in situ* grown on graphene oxide surface using polyethylenimine as reducing and stabilizing reagents, and then it was used as support for aptamer immobilization to detect human  $\alpha$ -thrombin from human serum.<sup>217</sup>

### 3.2.3. Use of nanoparticles coated with PEI as biosensors.

In many instances nanoparticles have been utilized in biosensor design, where the required amount of support is far lower than that in biocatalysis design. For example, glucose oxidase has been immobilized on nanotubes, using PEI to adsorb the enzyme and covalently immobilize ferrocene

derivatives to develop a glucose sensor (Figure 13).<sup>218</sup> Laccase has been immobilized on gold nanoparticles coated with PEI and used as biosensor.<sup>219</sup> Lactate oxidase was covalently immobilized on platinum nanoparticles supported on graphitized carbon nanofibers using PEI and glutaraldehyde (GA).<sup>220</sup>

### 3.3. Use of PEI to coat cell wall to co-immobilize enzymes and cells

Coimmobilization of enzymes and cells has significant interest. For example, glucose oxidase was immobilized on PEI-coated yeast cells and attached to cotton thread (Figure 14). The crosslinking with glutaraldehyde avoided undesired enzyme release and permitted several reuses without losing enzyme activity.<sup>221</sup>

In another paper, PEI was used to form PEI- D-lactate dehydrogenase - *Candida boidinii* aggregates that were later adsorbed onto cotton cloth.<sup>222</sup> That way, D-lactate dehydrogenase (LDH) was used to transform 2-oxo-4-phenylbutyric acid to R-2-hydroxy-4-phenylbutyric acid oxidizing NADH to NAD<sup>+</sup>, which was regenerated by the formate dehydrogenase presented in the cells of *Candida boidinii*.

### 3.4. Use of PEI to activate complex reactors to immobilize enzymes in the reactor itself

The immobilization of enzymes on complex reactors makes mounting and dismounting of the reaction complex (Figure 9). Thus, the use of some reversible immobilization enzyme protocol to prevent the necessity of reactor modification becomes a requirement. This might be exemplified in the use of monoliths (Figure 9), a new support configuration that is receiving increasing attention in biocatalysis<sup>223–229</sup> and chromatography.<sup>230,231</sup> Monoliths have better mass transfer hydrodynamics than

their standard particulate counterparts,<sup>232</sup> becoming very useful systems to treat high volumes of substrate. However they require a correct layout to ensure the correct circulation of the substrate, avoiding the promotion of preferential ways, considering that they are adequate when high substrate flow is utilized.

The activation of complex reactors with PEI is a good alternative for enzyme immobilization. Lactase from *Aspergillus oryzae* was immobilized on high-porosity mullite advanced ceramic material coated with PEI,<sup>233</sup> showing that the open microstructure of the monoliths permitted a good accessibility to the enzyme molecules immobilized inside the walls of the monolith and permitted high enzyme loadings. Later, classical cordierite and acicular mullite (ACM) monoliths were coated with silica and functionalized with PEI using different protocols.  $\beta$ -galactosidase from *Aspergillus oryzae* was then immobilized on them.<sup>234</sup> An extracellular lipase from *Yarrowia lipolytica* was one-step immobilized and purified from lipases from *Candida* sp. using poly(glycidyl methacrylate-co-ethylene glycol dimethacrylate) monolithic columns functionalized with PEI.<sup>235</sup> Four lipase isoforms from *Candida* sp. 99-125 were separated (isoform A, isoform B, isoform C and isoform D).

Acetylcholinesterase and choline oxidase were co-immobilized on a monolith and used for the evaluation of acetylcholinesterase inhibition caused by diverse agents.<sup>236</sup> Cryogels with high catalyst density were produced through cryostructuration of whole *Escherichia coli* cells (forming a monolith) containing  $\beta$ -glucosidase and crosslinking PEI and polyvinyl alcohol activated with glutaraldehyde.<sup>237</sup>

Biomimetic silicification was used to trigger enzyme immobilization on the surface of the graphene oxide nanosheets/FeOOH hydrogel for constructing monolithic biocatalytic reactors.<sup>238</sup> The graphene oxide adsorbs PEI and this polymer is used as the mineralization-inducing agent, forming silica on the graphene oxide surface. Here, penicillin G acylase was trapped simultaneously. This monolithic system displayed simple recyclability and high stabilities.

On-chip glucose oxidase and choline oxidase immobilized monolith micro-reactors were used to analyze the kinetic features of the enzymes.<sup>239</sup> The monolith was produced employing a sol-gel method, which was later functionalized using PEI. Polycarbonate microchannels supports were activated with PEI and used to immobilize alkaline phosphatase.<sup>240</sup> A glucose oxidase reactor constituted by a microchannel assembled in poly(methyl methacrylate), coated with PEI, and connected to an amperometric detector which was used to quantify glucose.<sup>241</sup> In another paper, the same group used different materials to immobilize glucose oxidase in microchannels using PEI activated with glutaraldehyde.<sup>242</sup> Later, microreactors with immobilized ascorbate oxidase, catalase, glutamate dehydrogenase and a coimmobilized glucose oxidase and horseradish peroxidase were immobilized using similar strategies.<sup>243</sup>

Similarly, PEI has been used to activate membranes to immobilize enzymes in membrane reactors. For example, lipase B from *C. antarctica* has been immobilized in an  $\alpha$ -alumina ceramic support activate with gelatin-PEI and successfully utilized in the production of butyl butyrate.<sup>244</sup>

PEI has also been employed to immobilize proteins in chips for ELISA assays even though these supports, as commented before, will have a strong capacity as anion exchangers. Trace levels of  $\alpha$ -fetoprotein (hepatocellular carcinoma biomarker) have been detected using a monoclonal antibody covalently immobilized on the amino groups of the PEI using poly(methyl methacrylate) microfluidic chips and electrochemical detection systems.<sup>245</sup> Enzyme microencapsulation (Figure 15) has been frequently used to immobilize enzymes, in many instances used later in a microreactor. For example, laccase was encapsulated via interfacial cross-linking of PEI and packed into a capillary-sized reactor with off-line capillary electrophoresis for substrate quantification.<sup>246</sup>

A mixture of lactate oxidase with PEI and poly(carbamoyl)sulphonate hydrogel was used for enzyme immobilization onto a platinum disk of a transducer in a system developed for development of L-lactate determinations.<sup>247</sup>

#### 4 Use of PEI as glue to immobilize multilayers of enzymes

##### 4.1. Use of PEI to form multilayers of the same enzyme

The formation of enzyme multilayers may permit increasing enzyme loading on a support (Figure 7). This may be interesting using porous supports, but becomes critical when the enzyme is immobilized on macroscopic non-porous supports having low specific area. The main point to be considered in order to succeed is to keep the enzyme activity of the in-between layers. To do this, enzyme distortion and diffusional problems need to be kept to the minimum. PEI, a random coil and flexible polymer, may be a very good alternative to act as glue between enzyme layers, or any other molecule. As previously discussed, the flexible structure permits to induce scarce alterations on the protein, and if a protein is coated with PEI and then another protein molecule is immobilized, this second protein will not interact with the first protein, but with PEI.

Glucose oxidase is one of the most widely used enzymes for this strategy. For example, PEI and poly(dimethyldiallyl- ammonium chloride) were assembled in alternate layers to achieve molecular films, without influence from substrate diffusion at up to 5  $\mu\text{g}$  of immobilized enzyme in 50  $\text{m}^2$ .<sup>248</sup> The enzyme film was also prepared by a premixing method, in which a glucose oxidase polyion complex was assembled alternately with another oppositely-charged polymer. This permitted to increase by 67-folds the activity compared to the conventionally assembled films. In another paper, multilayers of glucose oxidase using different ionic polymers as glue were produced onto a thiolated-gold surface and the resulting bioelectrode was used for glucose biosensing.<sup>249</sup> The supramolecular

structure was prepared by gathering polyethylenimine and Nafion (as anti-interference barrier), followed by adsorption of DNA (as stabilizing layer) and polyethylenimine and finally by the alternate deposition of polyethylenimine and glucose oxidase (as the biocatalytic film).<sup>249</sup> Glucose oxidase was also immobilized on multiwall carbon nanotube-modified glassy carbon electrode, using a layer of PEI to immobilize the enzyme. For multilayers, the next layer of PEI was directly assembled on the enzyme.<sup>250</sup>

Galactooligosaccharides were produced from lactose using  $\beta$ -galactosidase from *Aspergillus oryzae* immobilized on cotton cloth via PEI as biocatalyst - enzyme aggregate formation and growth of aggregates on individual fibrils of cotton cloth leading to multilayer immobilization of the enzyme.<sup>251</sup> In this case, 250 mg of enzyme per g of support could be immobilized. Later, a similar protocol was used using as support low-pressure plasma-modified cellulose acetate membrane.<sup>252</sup> Lactase was also covalently attached to low-density polyethylene for in-package production of lactose-free dairy products.<sup>253</sup> Multilayers of the enzyme were immobilized alternating polyethylenimine, glutaraldehyde and lactase (the 5-layer sample reaching up to  $1.3 \mu\text{g lactase}/\text{cm}^2$ ), although the affinity of the immobilized lactase towards the substrate remains unchanged after immobilization, the  $k_{\text{cat}}$  showed a decrease with each layer of attached lactase. In our opinion, this may be due to glutaraldehyde modification of the enzyme.

Urease has been immobilized on the walls of silicon microchannels by alternating layers of polycationic polymers (PEI, polydiallyldimethylammonium or polyallylamine combined) and polyanions (polystyrenesulfonate or polyvinylsulfate).<sup>254</sup> The best results were obtained using PEI/polystyrenesulfonate/PEI/urease/PEI (repeating this model for more layers). This strategy was used later by the same group to immobilize organophosphorous hydrolase on the same support.<sup>255</sup>

Single polycation cross-linked multilayers, based on polyethyleneimine, have been prepared using a method of 3,3',4,4'- benzophenonetetracarboxylic-dianhydride (BTCDA)-mediated

electrostatics and hydrogen bonds layer-by-layer assembly.<sup>256</sup> Different PEI polymers were adsorbed onto silica microparticles or silicon wafers, crosslinked using 3',4,4'- benzophenonetetracarboxylic dianhydride (that introduced carboxylic acids to the polymer), and used to adsorb a new layer of PEI. This was later used to immobilize pepsin or lysozyme using glutaraldehyde.

Lipase from *T. lanuginosus* was immobilized into multiple layers using polyethylenimine-treated cotton flannel cloth,<sup>257</sup> utilizing the enzyme property of forming bimolecular aggregates.<sup>258,259</sup> The use of aggregated enzyme permitted to increase the activity by 30 %.<sup>257</sup>

Nanostructures may also be glued with PEI forming multilayers. For example, microcylinders of halloysite of 50 nm diameter, 500 nm length and with 20 nm diameter hollow inner core were assembled by alternate adsorption with PEI, yielding ordered multilayers containing up to 20 layers of tubules connected by PEI.<sup>260</sup> The immobilization of alcohol dehydrogenase and halloysite loaded with NADH in alternation with PEI was achieved, providing a direct supply of NADH to the enzyme.

Thus, PEI has revealed itself as a powerful tool to produce many layers of enzymes keeping good activity even on the enzymes immobilized near the support surface.

#### 4.2. Use of PEI as glue to co-immobilize enzymes.

This strategy has been also used to co-immobilize two enzymes that may be coated with PEI and poly(styrenesulfonate). For example, layers of glucoamylase and glucose oxidase were assembled on an ultrafilter by layer-by-layer adsorption using PEI as glue.<sup>261</sup> That way, starch was converted to gluconic acid and H<sub>2</sub>O<sub>2</sub>. The optimal global activity was obtained when the glucoamylase layer was deposited on the layer of glucose oxidase.<sup>261</sup> In another paper, using alternating layers of PEI and poly(styrenesulfonate), a multilayer of alkaline phosphatase and glucose oxidase were formed,<sup>262</sup> although enzyme coimmobilization does not appear really advantageous in this case.<sup>23</sup>



In another research, a layer of poly(styrene sulfonate) was formed by adsorption on the surface of several supports (platinum electrode, glass plate, quartz crystal microbalance, quartz plates, mica and silicon substrates) followed by a layer of PEI.<sup>263</sup> This multilayer strategy was used to trap cholesterol esterase and cholesterol oxidase to detect cholesterol. In another paper, the enzymes lactate oxidase (to detect lactate) and ascorbate oxidase (to eliminate interferences with ascorbic acids) were co-immobilized on Pt electrodes with Ceria nanoparticles (as an oxygen deposit) using PEI as immobilizing agent and using several layers of each enzyme.<sup>264</sup> The final configuration of the biosensor was: Pt/CeO 2–3 layers of PEI/Lactate oxidase-3 layers of PEI/ascorbate oxidase.

In the examples listed above all enzymes were co-immobilized via ion exchange. That means that when the least stable enzyme is inactivated, both enzymes need to be discarded,<sup>23</sup> being this one of the frequently overlooked problems of enzymes coimmobilization. However, a strategy has been very recently proposed that involves the formation of layers of the involved enzymes using PEI as glue between one enzyme and the other, to prevent this coimmobilization drawback (Figure 16). The strategy was proposed to solve this problem in one specific situation, when one of the enzymes can be immobilized via ion exchange, and, at the same time, cannot be highly stabilized via multipoint covalent attachment. Moreover, this immobilized enzyme should be less stable than the other immobilized enzyme. The proposal was to immobilize a lipase via interfacial activation on octyl-agarose beads,<sup>31</sup> to coat the enzyme with PEI, and then, to immobilize the second enzyme on this PEI layer (Figure 16).<sup>265</sup> After inactivation of the enzyme, it could be desorbed and the lipase could remain immobilized and active on the support. It was found that enzyme desorption was not so simple, as the inactivated enzyme maximized the enzyme-polymer interactions.<sup>120,121</sup> As some PEI was desorbed when the second enzyme was released, the immobilized enzyme required to be incubated with PEI before immobilizing a new batch of the second enzyme. This problem is not relevant at laboratory

scale but it is quite relevant at industrial level. It has been partially solved using octyl-glyoxyl agarose and modification of the lipase with glutaraldehyde.<sup>266</sup>

## 5. Use of PEI to enhance the formation of solid supports formed by ionic polymers

There are many examples in the literature where PEI is not used to activate supports, but to improve the physical stability of supports formed by interaction between anionic polymers and divalent cations, mainly to trap enzymes. Trapping of enzymes should not improve enzyme rigidity, but may generate an adequate enzyme environment or prevent subunit dissociation of multimeric enzymes.

One example is the case of supports formed by alginate or  $\kappa$ -carrageenan and calcium.<sup>267</sup> Although most examples use chitosan to reach this goal,<sup>268</sup> some examples may be found using PEI to stabilize this kind of supports and used to immobilize D-hydantoinase<sup>269</sup> or penicillin acylase.<sup>270</sup> Moreover this modification allows introducing glutaraldehyde groups on the support (via the primary amino groups of PEI), and prevents enzyme release, a usual problem using trapping techniques for enzyme immobilization.<sup>271</sup> Some examples may be found in the literature of this use of PEI, for example the immobilization of lipase from *Expansum penicillium* to be used in the esterification of 1-dodecanol with dodecanoic acid in benzene,<sup>272</sup> porcine pancreatic lipase to hydrolyze olive oil,<sup>273</sup>  $\beta$ -glucosidase from *Aspergillus niger* to produce reducing sugar from algae cellulosic residue.<sup>274</sup> Calcium pectinate beads were treated with PEI and glutaraldehyde and used to immobilize  $\beta$ -D-galactosidase.<sup>275</sup>

In other instances, PEI is utilized as nucleation point to get a solid support. Several enzymes have been trapped by silica co-precipitation. The process consists in a mixture of enzyme and PEI, and the further incorporation of a phosphate-buffered silicate solution. This formed a colloidal co-

precipitate consisting of the enzyme and PEI within a hydrated, amorphous silica matrix.<sup>276</sup> Later a similar strategy was used to coat the surface of the immobilization support with silicic acid, and later used to immobilize several proteins via ion exchange.<sup>277</sup>

In another paper, N-isopropylacrylamide, N,N-methylene bis-acrylamide and PEI were utilized to form a microgel via adsorption.<sup>278</sup> This microgel was used to immobilize acetyl CoA synthetase, and finally the microgel was covalently immobilized upon polyethylene terephthalate track etched membrane.<sup>278</sup>

Other research shows a method based on polyphenol chemistry for the facile preparation of microcapsules in four steps: production of a sacrificial template, assembly of the polyphenol coating on the template surface, cross-linking of the polyphenol coating by cationic polymers (PEI), and removal of the template.<sup>279</sup> The strong interfacial affinity of tannic acid helped towards the generation of polyphenol coating through oxidative oligomerization, while the reactivity of tannic acid permitted the cross-linking with PEI through Schiff base/Michael addition reactions. Glucose oxidase was immobilized in the microcapsules, showing improved pH and thermal stabilities and good catalytic activity.<sup>279</sup>

## 6. Use of PEI capsules to immobilize enzymes

Encapsulation of enzymes inside PEI capsules has been described as a very efficient way to avoid dissociation of multimeric enzymes (Figure 3). For example, glutamate dehydrogenase from *Thermus thermophilus* and formate dehydrogenase from *Pseudomonas* sp. were coated with PEI and treated with glutaraldehyde to prevent enzyme dissociation under acidic pH value.<sup>188</sup> This permitted to have a free enzyme where dissociation was not possible, enlarging the range of conditions where the enzyme may be handled, in operation or just to later immobilize the composite.

The enzyme-PEI composite was ionically exchanged on cationic exchangers. For example, formate dehydrogenase could be treated with PEI and glutaraldehyde after adsorption on cationic exchangers, where the native enzyme cannot become immobilized. Later, a similar strategy was used to stabilize glutamate dehydrogenase from *Escherichia coli*, showing that together with preventing enzyme dissociation, this strategy also improved enzyme stability in stirred systems.<sup>189</sup>

Glucose oxidase was coated with PEI and this stabilized enzyme/polymer composite was immobilized into woodceramics that served as both immobilization matrix and electrochemical transducer.<sup>280</sup> Again, immobilization was only possible after coating the enzyme with PEI to generate a cationic enzyme surface. A cross-linked PEI wall was generated from an emulsion of an aqueous phase containing laccase, forming microcapsules that were later immobilized on the surface of the biosensor electrode.<sup>281,282</sup> The laminar jet break-up technique was used to generate PEI microcapsules for immobilization of glucose oxidase from *Aspergillus niger* and laccase from *Trametes versicolor*.<sup>283</sup> Later, these enzyme/PEI microcapsules were used to immobilize the enzymes in paper.<sup>284</sup> In another research effort, ink was supplemented with microcapsules produced via interfacial polycondensation of PEI with the sebacoyl chloride and containing laccase from *Trametes hirsuta* or *Trametes versicolor*.<sup>285</sup> Then, they were printed or coated on a paper substrate.

In another paper, an aqueous solution of PEI was mixed with an aqueous solution of Fmoc-diphenylalanine peptide and sodium silicate.<sup>286</sup> This way a new class of organic-inorganic hybrid capsules (FPSi) with multi-layered structure was fabricated and used to encapsulate epoxide hydrolase and bovine serum albumin.

## 7. Use of PEI in the preparation of crosslinked enzyme aggregates (CLEAs)

The immobilization of enzymes by CLEAs technology was proposed by Prof Sheldon's group.<sup>287,288</sup> This protocol was a simplification of the enzyme crosslinked crystals,<sup>289,290</sup> consisting in

the precipitation of the enzymes (adding some precipitant agent) and their further crosslinking (usually with glutaraldehyde, although in some instances some alternatives have been used).<sup>291,292</sup> That way the enzyme aggregate cannot solubilize when the precipitant agent is removed (Figure 17). Advantages involve that enzyme purity is not a requirement, and aggregation is far simpler than crystallization. Thus, combi-CLEAs have been produced by co-aggregating several enzymes.<sup>293–295</sup> However, in some instances the crosslinking step may become a problem if the target protein is poor in Lys groups in the surface. This may be solved using bovine serum albumin or other protein feeder,<sup>296–298</sup> or if the enzyme could be chemically aminated.<sup>299</sup> As all the solid is enzyme, diffusion problems tend to be high.<sup>23</sup> this may be partially reduced using an inert protein.<sup>300</sup>

PEI may play a significant role in facilitating the crosslinking of aggregates, as it is very rich in primary amino groups, and also reduces the protein density decreasing substrate diffusional limitations. Thus, stable CLEAs of glutaryl acylase,<sup>301</sup> lipases from *Alcaligenes sp.*, *C. antarctica* (fraction B), *Geotrichum sp.* and *Serratia marcescens*<sup>302–304</sup> could be prepared using PEI as a feeder while the direct crosslinking failed. In the case of lipases, enzyme features (e.g., specificity) could be also tuned using PEI.<sup>302</sup>

The cationic nature of PEI has been used to generate “saline” environments (Figure 6), and has improved the stability of CLEAs of oxygen-labile nitrilases versus oxygen<sup>305</sup> or penicillin G acylase versus organic solvents (mixing PEI and dextran sulfate).<sup>306</sup> These penicillin acylase preparations were used in the synthesis of diverse antibiotics.<sup>307–309</sup>

## 8. Modification of immobilized enzymes with PEI

The points above showed examples where PEI was used to activate the support or to form the own support. Now, we will give a view on the employment of PEI to modify previously immobilized molecules to improve different features, from stability to specificity or activity.

### 8.1. Intersubunit crosslinking of multimeric enzymes by PEI immobilization or PEI modification of immobilized enzymes

The first step in the inactivation of multimeric enzymes involves in many instances dissociation of the enzyme.<sup>310–312</sup> The simultaneous immobilization of all enzyme subunits on a support prevents this inactivation cause (Figures 2 and 3). However in some instances the geometry of the enzyme prevents involving all enzymes in the immobilization.<sup>24</sup> For these instances, the coating of the enzyme with a polymer has shown to be a powerful tool to crosslink all enzyme subunits, although in many instances aldehyde dextran has been utilized.<sup>313,314</sup> In fact, in some instances enzymes immobilized on PEI coated supports have been further stabilized by treatment with aldehyde dextran. Nucleoside 2'-deoxyribosyltransferase *Bacillus psychrosaccharolyticus* was immobilized on PEI coated agarose beads followed by crosslinking with 70% oxidized dextran.<sup>147</sup> This biocatalyst was later used to produce trifluridine.<sup>146</sup> A similar strategy was utilized with a pyrimidine nucleoside phosphorylase from *Bacillus subtilis* and a thymidine phosphorylase from *Escherichia coli*.<sup>315</sup> The biocatalysts were used in the synthesis of 5-halogenated pyrimidine 2'-deoxyribonucleosides by transglycosylation. Nucleoside phosphorylases from *Citrobacter koseri*, *Clostridium perfringens* and *Streptococcus pyogenes* were also immobilized on PEI coated supports and crosslinked with partially oxidized aldehyde dextran.<sup>316</sup>

The homodimeric thymidine phosphorylase from *Escherichia coli* was immobilized via ion exchange on PEI coated agarose and Sepabeads ® and cross-linked with partially oxidized dextran

aldehyde to prevent enzyme dissociation.<sup>317</sup> These biocatalysts were employed for the one-pot synthesis of 5-fluoro-2'-deoxyuridine starting from 2'-deoxyuridine and 5-fluorouracil. Uridine phosphorylase from *Bacillus subtilis* was immobilized/stabilized onto Sepabeads coated with polyethyleneimine and cross-linked with aldehyde dextran (UP-Sep-PEI-Dx).<sup>145</sup> This biocatalyst was used in combination with immobilized purine nucleoside phosphorylases in the synthesis of 2' - deoxyguanosine by enzymatic transglycosylation in aqueous solution between 2'-deoxyuridine and guanine.<sup>318</sup>

In another example, lipase dimers were immobilized on PEI-coated supports. Lipases have a certain tendency to form bimolecular aggregates confronting the open form of two lipase molecules with clearly altered functional properties, and their immobilization in PEI coated supports seem to keep these dimers.<sup>258</sup>

However, there are a handful of examples of the use of PEI as the final crosslinking agent. For example, the tetrameric  $\beta$ -xylosidase from *Selenomonas ruminantium* was immobilized on glyoxyl agarose beads, with a slight stabilization due to the non-implication of all enzyme subunits.<sup>319</sup> The modification of the immobilized enzyme with aldehyde-dextran or polyethylenimine fully stabilized the quaternary structure of the enzyme and greatly improved enzyme stability. Another paper shows that the immobilization of the multimeric alcohol dehydrogenase from baker's yeast on agarose activated with glyoxyl groups stabilized the enzyme by 50-folds.<sup>320</sup> However, not all enzyme subunits become immobilized. Physical modification of the immobilized enzyme with PEI permitted to crosslink all enzyme subunits, improving enzyme stability and preventing enzyme dissociation.

In another paper, the heterodimeric cephalosporin C acylase was ionically exchanged in the aminated support LX1000-HA, treated with glutaraldehyde and later treated with PEI to improve the enzyme stability.<sup>321</sup>

Thus, the large size of PEI may be a very useful way to physically crosslink multimeric enzymes. However, it may be expected that this protection is only shown under conditions where PEI remains interacting with the enzyme (that is, it may be lost at high ionic strength or extreme pH values). Treatment of the modified enzymes with glutaraldehyde or other bifunctional reagents may make this strategy useful under all conditions, but it is not usually performed.

## 8.2. Generation of hydrophilic environments by PEI coating

The generation of hydrophilic environments (Figure 4) may be used to partition some detrimental hydrophobic compounds from the enzyme environment (Figure 6) and thus, to improve enzyme performance.

A general procedure for stabilization of O<sub>2</sub>-labile enzymes exploiting "salting out" of oxygen from highly hydrophilic enzyme microenvironment was proposed by Klivanov's group a long time ago, suggesting that PEI could be a very suitable support activating agent for this goal.<sup>69</sup>

The generation of highly hydrophilic environments surrounding enzyme molecules has been proposed as a strategy to produce a partition of hydrophobic organic solvents and that way reduce their concentration in the enzyme surrounding (Figures 4 and 6). This has been applied to achieve penicillin G acylase immobilized biocatalyst that coupled a high rigidification by immobilization via multipoint covalent attachment to a highly hydrophilic environment formed by "solid salts" formed by aldehyde dextran and PEI-dextran sulfate layers, and that were highly stabilized in the presence of organic solvents.<sup>322,323</sup> These biocatalysts could be used in the hydrolysis of penicillin G in the presence of organic solvents<sup>324</sup> in the thermodynamically controlled synthesis of antibiotics,<sup>309,325</sup> or other amides,<sup>326</sup> or in the enantioselective synthesis of phenylacetamides.<sup>327</sup>

The coimmobilization of penicillin G acylase and PEI did not improve the stability of the immobilized enzyme. This was attributed to the fact that the enzyme was not adsorbed on PEI coated



supports,<sup>328</sup> therefore the polymer did not really prevent the exposition of enzyme to high organic solvent concentrations due to the lack of enzyme-polymer interactions. The genetic introduction of eight additional Glu residues homogenously distributed throughout the enzyme surface permitted the polymer adsorption, and thus enzyme stabilization versus organic solvents by coimmobilization of the enzyme and the polymer.<sup>329</sup>

In another paper, lipases from *Rhizomucor miehei* and *Candida rugosa* were immobilized on Sepabeads decaoctyl and coated with PEI to decrease the interactions between the immobilized lipase molecules and the organic solvents when using these derivatives in anhydrous media.<sup>330</sup> These biocatalysts were more active and stable than the uncoated preparations when catalyzing esterifications and transesterifications in anhydrous media. In another instance, the coating of a commercial preparation of lipase B from *C. antarctica* (form B) (Novozym 435) with PEI improved their stability in the production of structured lipids, mainly by preventing the formation of crystals of aggregated material and improving the support stability.<sup>331,332</sup> This use of PEI has been scarcely explored although it is a simple way to protect enzymes from hydrophobic compounds.

### 8.3. Physical crosslinking of reversibly immobilized enzymes

Physical immobilization of enzymes is very popular, because it is very simple to implement, the supports tend to be stable, and the immobilization is reversible, enabling the reuse of the support when the enzyme has been inactivated. This immobilization is usually not very efficient in improving enzyme stability, except when multimeric enzymes are involved. However, there is a case where the physical immobilization produces a high stabilization of the enzyme: the immobilization of lipases on hydrophobic supports.<sup>333</sup> This stabilization is even higher than using multipoint covalent immobilization,<sup>32,33</sup> because the lipase interfacially activated versus the hydrophobic surface of the

supports gives a lipase form that is more stable than the enzyme in closed/open conformation, even if this form is stabilized via multipoint covalent attachment<sup>34,334,335</sup>

However, lipases are desorbed from the hydrophobic support under drastic conditions,<sup>336</sup> or even by the action of some substrates with detergent properties<sup>35,36,337</sup> producing product contamination by the enzyme and biocatalyst inactivation. This has been solved using heterofunctional supports combining acyl moieties to achieve interfacial activation of the lipase and covalent reactive groups.<sup>336,338–342</sup> Also, covalent crosslinking has been performed using aldehyde dextran.<sup>343</sup> A simpler solution has been recently proposed: the coating of the fully loaded enzyme preparations with PEI and dextran sulfate greatly reduced enzyme desorption and significantly improved enzyme stability (Figure 18). Thus, some lipases immobilized on octyl agarose, like those from *Rhizomucor miehei*,<sup>344</sup> *C. antarctica* (form B),<sup>345,346</sup> *T. lanuginosus* or the phospholipase Lecitase have been stabilized by coating with PEI.<sup>347</sup> That is, coating of octyl-lipase preparations with PEI permits to reduce enzyme leakage without losing the reversibility of the immobilization of the lipases, and the positive effects of stabilization versus hydrophobic molecules may be also achieved.

#### 8.4. Bioimprinting of enzyme features

Enzyme bioimprinting may be defined as a strategy to maintain an enzyme conformation induced by a given agent in the absence of this particular agent.<sup>348–350</sup>

PEI or supports coated with PEI have proven to be very efficient for this goal. For example, the activity of Lecitase Ultra covalently immobilized on cyanogen bromide was increased if incubated in the presence of some detergents.<sup>351</sup> However, the detergents were negative for enzyme stability.<sup>351</sup> To keep this hyperactivation, which was associated to the stabilization of the open form of the enzyme, the immobilized enzyme was incubated with PEI in presence of sodium dodecyl sulfate, achieving a

Lecitase form 50-folds more active than the initial preparation. It was showed that the PEI treatment maintained the open form of the lipase in absence of the detergent (Figure 19).<sup>351</sup>

Immobilization of lipase from *C. antarctica* (form B) on PEI supports at different pH values permitted to keep the properties exhibited by the enzyme under the immobilization pH values in the resolution of R,S-mandelic acid methyl ester, even though the reaction was performed at other pH values.<sup>352</sup>

Finally, the oligomeric state of invertase from *S. cerevisiae* depends on the pH, presenting different properties.<sup>353,354</sup> The enzyme was immobilized on Sepabeads coated with PEI at different pH values where the aggregation of the enzyme was different. These biocatalysts presented different stabilities, suggesting that the immobilized enzyme kept this aggregation state under any pH value.<sup>355</sup>

### 8.5. Modification of enzyme properties

It has been reported how different immobilization strategies or the further chemical or physical modification may be used to tune enzyme selectivity, specificity or activity.<sup>22,23,45,46</sup> In many examples, PEI-coated supports or modification with PEI of immobilized preparations (Figure 5) have been used to achieve this alteration in enzyme features.

In some cases, PEI was used to modify the properties of immobilized enzymes. Lecitase immobilized in cyanogen bromide agarose or on octyl-agarose and further coated with PEI permitted to significantly increase enzyme activity (even by 30 folds using the octyl-Lecitase-PEI at pH 5 in the hydrolysis of methyl phenyl acetate).<sup>356</sup> Lipases from *C. antarctica* (form B), *T. lanuginose* and *Rhizomucor miehei* were immobilized on CNBr-activated Sepharose beads or onto octyl-agarose and used in the hydrolysis of sardine oil and submitted to different chemical or physical modifications,

among them with PEI.<sup>357</sup> The selectivity and activity of lipase from *T. lanuginosa* immobilized on octyl-agarose increased two-fold after PEI modification.

In other cases, immobilization of the lipases on PEI-coated supports enables the alteration of their catalytic properties. Lipase B from *C. antarctica* was immobilized on different supports and used in the hydrolytic resolution of several ( $\pm$ )- $\alpha$ -hydroxy-phenylacetic acid derivatives.<sup>358</sup> The enzyme immobilized on cyanogen bromide presented an enantiospecificity of 7.4, while the enzyme immobilized in PEI coated agarose exhibited an enantiospecificity of 67 in the hydrolysis of  $\alpha$ -hydroxy-phenylacetic acid methyl ester under similar conditions. Moreover, the enantioselectivity of the PEI derivative decreased from 67 to 14 when the temperature went from 25 to 4°C at pH 5, while the E of some other derivatives improved significantly under similar experimental changes.<sup>358</sup>

The enantiospecificity of the lipase from *Candida rugosa* in hydrolysis of R,S methyl mandelate went from <2 using some preparation to around 200 for the towards S-isomer using a lipase immobilized on PEI coated support.<sup>359</sup>

The lipase from *Aspergillus niger* immobilized on PEI coated support was six times more active than a covalent preparation in the hydrolysis of 1-thioisopropyl-2,3,4,6-tetra-O-acetyl- $\beta$ -d-galactopyranoside. However, this covalent preparation was more active in the hydrolysis of galactal.<sup>360</sup> The lipase from *Candida rugosa* immobilized on octyl agarose was completely specific and regioselective in the hydrolysis of galactal, producing the C-6 monohydroxylated product in 99% yield while the enzyme immobilized on PEI coated agarose beads hydrolyzed in C-6 but also in C-3 positions.<sup>360</sup> Lipase from *Aspergillus niger* immobilized on supports coated with PEI was 800 times more active than the enzyme immobilized on octyl-agarose in the hydrolysis of 2-acetamido-2-deoxy-1-(4-nitrophenyl)-3,4,6-tri-O-acetyl- $\beta$ -d-glucopyranoside.<sup>361</sup> However, the octyl derivative was five times more active than the PEI preparation in the hydrolysis of 1-(4-nitrophenyl)-2,3,4-tri-O-acetyl- $\beta$ -d-xylopyranoside.

Lecitase immobilized on PEI coated supports was much more active in the hydrolysis of 1,2,3,4,6-penta-*O*-acetyl- $\alpha$ -d-mannopyranose or 2-acetamido-2-deoxy-1,3,4,6-tetra-*O*-acetyl- $\beta$ -D-glucopyranose than the enzyme immobilized on octyl-agarose, but it was 4 folds less active in the hydrolysis of 1,2,3,4,6-penta-*O*-acetyl- $\beta$ -d-galactopyranose.<sup>362</sup>

Among the different enzymes and immobilized biocatalysts, lipase from *T. lanuginosus* immobilized on supports coated with PEI was the most efficient catalyzing the hydrolysis of 1,2,3,4,6-penta-*O*-acetyl- $\beta$ -D-glucopyranose, permitting us to obtain up to 70% of the 6-hydroxy product.<sup>363</sup> In the hydrolysis of 2-acetamido-2-deoxy-1,3,4,6-tetra-*O*-acetyl- $\beta$ -D-glucopyranose, the form B of the lipase from *C. antarctica* immobilized on octyl agarose was not regioselective. However, the enzyme immobilized on supports coated with PEI was highly specific and regioselective producing the 6-hydroxy-2-acetamido-2-deoxy-1,3,4-tri-*O*-acetyl- $\beta$ -D-glucopyranose in 70 % yield.<sup>363</sup> A lipase from porcine pancreas was adsorbed on PEI-coated support and utilized in the resolution of ( $\pm$ )-glycidyl butyrate, showing an enantiospecificity of 6.<sup>127</sup> The treatment of the adsorbed lipase with glutaraldehyde permitted to increase this value to E = 61. The use of lipase B from *C. antarctica* immobilized on a support coated with PEI permitted to obtain (S)-glycidyl butyrate (90% ee), while other immobilized preparations offered much lower values.<sup>364</sup>

A contaminant esterase contained in commercial extracts of *C. antarctica* (form A) was immobilized on supports coated with PEI and used in the enantioselective hydrolysis of (3*RS*,4*RS*)-trans-4-(4'-fluorophenyl)-6-oxo-piperidin-3-ethyl carboxylate hydrolyzing mainly the (3*S*,4*R*) isomer with an enantiospecificity higher than >100.<sup>365</sup> Lipase from *Candida rugosa* immobilized on PEI coated supports was the most active hydrolysing 2-phenyl-2-butyroylacetic acid at pH 7 or methyl mandelate.<sup>366</sup>

## 8.6. Use of PEI to co-immobilize cofactors, conductive molecules and enzymes

In biosensor design, a good conductivity between enzyme and electrode is a critical requirement for good performance.<sup>367</sup> In this sense the modification of PEI with different molecules and their coimmobilization with the enzyme has been a successful alternative (Figure 13). For example, glucose oxidase coated with PEI was immobilized on polypyrrole film deposited on poly(styrene sulfonate) modified glass, mica and indium-tin-oxide coated glass plates.<sup>368</sup> The immobilization of glucose oxidase on a conductive polypyrrole film permitted to build glucose biosensors. In another paper, PEI was adsorbed on the surface of a screen-printed carbon electrode and ferricyanide ions were adsorbed and confined inside the polymer bed.<sup>369</sup> This was incubated in a  $\text{FeSO}_4$  solution, to permit  $\text{Fe}^{2+}$  cations to react with the adsorbed  $[\text{Fe}(\text{CN})_6]^{3-}$ . The reaction led to the formation of Prussian blue crystals that were very active in the electrochemical reduction of hydrogen peroxide. After glucose oxidase was immobilized via ion exchange on the electrode surface, a biosensor for glucose was produced. This sensor reduced the redox activity of many interfering compounds (Figure 13).<sup>369</sup> PEI was modified with ferrocene groups and the modified polymer was adsorbed on multi-wall carbon nanotubes attached carbon cloth.<sup>370</sup> Then, it was crosslinked with glutaraldehyde and glucose oxidase was covalently immobilized onto this electrode.

A step beyond in the design of biosensors was the modification of PEI with ferrocene and  $\text{NAD}^+$ . This polymer was coimmobilized in an electrode with the enzyme alcohol dehydrogenase to build a reagentless alcohol biosensor.<sup>371</sup> A similar strategy was also used to coimmobilize PEI-ferrocene/ $\text{NAD}^+$  polymer<sup>and</sup> L-lysine 6-dehydrogenase from *Geobacillus stearothermophilus* in a gold sensor to build a reagentless L-lysine sensor. Several layers of enzyme and polymers were used to increase the sensor sensibility. PEI was covalently modified with  $\text{NAD}^+$  and 3,4-dihydroxybenzaldehyde.<sup>372</sup> The polymer displayed both electrochemical properties of catechol (useful as electron mediator) and biological activity of NAD moieties when using alcohol dehydrogenase. Just a slight increase in the Michaelis-Menten constant was observed with the polymeric  $\text{NAD}^+$ . The

bioelectrochemical properties of this double modified polymer were used to develop a reagentless biosensor using glucose-6-phosphate dehydrogenase. The polymer and the enzyme were retained on the electrode surface by using an ultrafiltration membrane.<sup>372</sup>

Nowadays the use of enzymes that require cofactors (NAD(P)H/NAD(P), ATP, UDP) is one of the aims of biocatalysis, as not only the cofactor regeneration but also its reuse for several reaction cycles may determine the implementation of many processes.<sup>373</sup> The immobilization of the cofactors on polymers is one of the preferred ways to facilitate this reuse, and PEI is one of the polymers used (as shown in the design of biosensors above). Related to biocatalysis, poly(allylamine) was modified with NADH and trapped with several dehydrogenases in hollow-fiber and used to produce methanol from CO<sub>2</sub>.<sup>374</sup>

In a very recent paper, López-Gallego and coworkers have shown that phosphorylated cofactors (PLP, FAD<sup>+</sup>, and NAD<sup>+</sup>) become “reversibly immobilized” by using supports coated with PEI (Figure 20).<sup>375</sup> It is postulated that the cofactors are retained in the polymer in a reversible way by ion exchange, facilitating the cofactor availability for the enzyme, and with very low levels of leaching. Two examples were given. In the first one, alcohol dehydrogenase from *Thermus thermophilus* was immobilized on glyoxyl agarose and PEI was coimmobilized on this support. As regenerating enzyme, they employed the formate dehydrogenase from *Candida boidinii*, which failed in becoming immobilized on glyoxyl agarose in an active form. For this reason, the enzyme was adsorbed on PEI treated with 1,4-butanediol diglycidyl ether to get a covalent immobilization of this enzyme on the PEI layer. Then, the combi-biocatalyst was incubated in a NADH solution and the biocatalyst with the adsorbed cofactor was used in the transformation of 2,2,2-trifluoroacetophenone to *S*-(trifluoromethyl)benzylalcohol, just using formic acid to regenerate the cofactor by formate dehydrogenase. The cofactor was re-cycled for 107 h of use (equivalent to 356 operational volumes) without significant NAD<sup>+</sup> losses. In another example, racemic methylbenzylamine was transformed in



*R*-methylbencylamine through S-selective deamination using w-transaminase coimmobilized in glyoxyl agarose with PEI and later incubated with PPL.<sup>375</sup> This paper is of outstanding importance in the design of reactions involving phosphorylated cofactors, and cofactor may be not only recycled but even reused several cycles of reaction. After the cofactor inactivation, if the enzyme remains active, although the authors did not study that, it is very likely that incubation at high ionic strength may release the inactive cofactor and permit the loading of fresh cofactor batch, reusing the coimmobilized enzymes.

## 6. Conclusions and future trends

From the ideas and results presented in this review, it seems obvious that the impact of PEI in the future may be even greater than the already outstanding one that we are currently experiencing. Its price is very competitive, the handling is simple, and its possibilities of use in biocatalyst design are huge. It is true that some proteins may be readily inactivated by PEI, but they are more the exception than they are the rule.

To take full advantage of these possibilities, a proper activation of the support (having a thick polymeric bed) and a thorough control of the immobilization conditions (to permit the penetration of the enzyme on the bed) are mandatory to have an adsorption involving the maximum percentage of the enzyme surface. This will permit to take full advantage of the use of this polymer, reaching a strong enzyme adsorption and also a higher protection of the enzyme from inactivating agents. Among the challenges that may limit the utilization of PEI in the design of biocatalyst is the possibility of release of the enzyme if the PEI is used to coat the support, or of the enzyme if the enzyme is coated with PEI. This will produce undesired effects and a contamination of the medium. One general solution may be to use enzymes with a surface greatly enriched in anion groups, and if possible, with a reduction in cationic groups. This may be achieved via chemical or genetic modification, and may help to solve this



problem. Very recently, it has been reported a possibility to prevent PEI desorption from PEI coated immobilized: the chemical immobilization of the PEI in enzymes surfaces activated with bifunctional reagents, like glutaraldehyde. This fully prevents this problem under any condition and may be used to coimmobilized enzymes or enzymes and cofactors.

Other drawback is also an advantage: the polymer flexibility. This flexibility avoids the generation of strong diffusion problems (that is positive), but also did not introduce rigidity to the enzyme, that will be that way hardly rigidified. This makes that PEI is not valid to immobilize or even intramolecular crosslink enzymes when the objective is increase enzyme rigidity to enhance enzyme stability. However, it may be very adequate when stability is not a problem, because the enzyme is already very stable in the operation conditions (e.g., thermophilic enzymes). Otherwise, enzyme rigidification should be accomplished by other genetic or chemical strategies.

This polymer flexibility also produces a further inconvenience: enzymes that may become inactivated by interactions in partially hidden pockets may be inactivated using PEI coated supports and not standard aminated supports. Branched PEI may limit this effect, if this may be controlled.

Moreover, this polymer is multifunctional, and it may permit partial modifications to tailor the final properties, or to introduce some required groups. This application is underutilized in the current literature and may become one of the greatest advantages of using a multifunctional polymer like PEI.

Some possibilities related to PEI are clearly infra-utilized, for example the use of PEI to produce multi-layers of enzymes on surfaces where the specific area is small. This may permit to dilute the enzyme with a random coil polymer (reducing mass transfer limitations) and to introduce high amounts of protein per surface unit. Even if some substrate diffusion limitations occur, this may not be detrimental for certain applications, e.g., it may enlarge the range of concentrations where a biosensor may increase the response when the concentration of the analyte is high. Some likely advantages of

PEI still remain unexplored. One point that (surprisingly) has not been studied at all is the strong buffering properties of PEI. We have not found any application of this property, but at first glance it may be enough to reduce or even fully prevent some pH gradients formed in enzyme reactions releasing basic or acids compounds, and it may also improve enzyme stability in titration processes by preventing enzyme exposition to drastic pH values.

Some others features are used, but have been very recently described and a wider diffusion of these characteristic is expected in the near future. Stabilization of multimeric enzymes, even immobilization of multi-enzymatic complexes is another application that should have more impact due to the simplicity of the protocols and efficiency of the treatments. Similarly, the physical crosslinking of physically immobilized enzyme molecules and the generation of enzyme nanoenvironments opens up new possibilities to solve some problems of very often used biocatalysts, approaching these biocatalysts to the industrial implementation for new processes. Prevention of enzyme inactivation due to interaction with hydrophobic interfaces (e.g., gas bubbles, drops of insoluble solvents) will have a more relevant role when the use of non-porous magnetic nanoparticles to immobilize proteins has a wider application.

However, the combination of two ideas very recently presented may be the main future application of PEI in biocatalysis. This is the involvement of PEI in cascade enzymatic reactions involving phosphorylated cofactors and several enzymes (Figure 21). Many enzymes involved in these reactions are multimeric ones that may be stabilized just via ion exchange in a polymeric bead involving all enzyme subunits, but in some cases they cannot be stabilized by multipoint covalent attachment for different reasons. Thus, among the enzymes involved in the cascade reaction, enzymes that are more stable or that can be significantly stabilized via multipoint covalent attachment may be immobilized following optimal protocols to improve at the maximum the enzyme properties. Later, they can be coated with PEI, which may present many positive effects on the immobilized enzyme

stability. If desired, it is possible to covalently attach PEI to the enzyme or to the support to prevent PEI desorption. Next, the second enzyme may be ionically exchanged on this polymer-coated enzyme, and finally, the combi-catalyst-PEI can be incubated in the solutions of the cofactors. That way, the cofactor may be reused for several cycles, and the least stable enzyme discarded after inactivation enabling the reuse of the most stable enzyme. The possibility of modifying PEI with different reagents may permit to facilitate electron transmission, or also enable the second enzyme to be immobilized via another reversible mechanism (e.g., thiol exchange). Genetic or chemical enrichment of the enzyme surface on anionic groups can make stronger second enzyme ion exchange if necessary. Combi-CLEAs of different enzymes and PEI may be incubated in cofactor solutions with similar objectives, although in this case the reuse of the most stable enzyme will not be possible.

Thus, the use of PEI in biocatalysis may be considered that still is in the first steps even though it has been produced for a long time. The road ahead may lead to solutions for biocatalysts design problems that may be solely in our imagination, or even situations which we have not even dared to dream yet.

## Acknowledgments

This work was supported by grants from Spanish Ministry of Economy and Competitiveness (MINECO) project number CTQ2013-41507-R, CNPq (process 403505/2013-5). A.B.M. thanks MINECO, Generalitat Valenciana and FEDER (CTQ2015-66080-R MINECO/FEDER and PROMETEOII/2014/010) for financial support.

972 **References**

- 973 1 P. Anastas and N. Eghbali, *Chem. Soc. Rev.*, 2010, **39**, 301–312.
- 974 2 M. Poliakoff, J. M. Fitzpatrick, T. R. Farren and P. T. Anastas, *Science (80-. )*, 2002, **297**, 807–  
975 810.
- 976 3 P. Schwager, N. Decker and I. Kaltenegger, *Curr. Opin. Green Sustain. Chem.*, 2016, **1**, 18–21.
- 977 4 V. S. Ferreira-Leitão, M. C. Cammarota, E. C. G. Aguiéiras, L. R. V. de Sá, R. Fernandez-  
978 Lafuente and D. M. G. Freire, *Catalysts*, 2017, **7**, 9.
- 979 5 A. Schmid, J. S. Dordick, B. Hauer, A. Kiener, M. Wubbolts and B. Witholt, *Nature*, 2001, **409**,  
980 258–268.
- 981 6 H. E. Schoemaker, *Science (80-. )*, 2003, **299**, 1694–1697.
- 982 7 D. J. Pollard and J. M. Woodley, *Trends Biotechnol.*, 2007, **25**, 66–73.
- 983 8 J. M. Woodley, *Trends Biotechnol.*, 2008, **26**, 321–327.
- 984 9 P. V Iyer and L. Ananthanarayan, *Process Biochem.*, 2008, **43**, 1019–1032.
- 985 10 V. G. H. Eijssink, S. Gåseidnes, T. V. Borchert and B. van den Burg, *Biomol. Eng.*, 2005, **22**,  
986 21–30.
- 987 11 K. M. Polizzi, A. S. Bommarius, J. M. Broering and J. F. Chaparro-Riggers, *Curr. Opin. Chem.*  
988 *Biol.*, 2007, **11**, 220–225.
- 989 12 V. V. Mozhaev and K. Martinek, *Enzyme Microb. Technol.*, 1984, **6**, 50–59.
- 990 13 D. Cowan, Q. Meyer, W. Stafford, S. Muyanga, R. Cameron and P. Wittwer, *Trends*  
991 *Biotechnol.*, 2005, **23**, 321–329.
- 992 14 T. Uchiyama and K. Miyazaki, *Curr. Opin. Biotechnol.*, 2009, **20**, 616–622.
- 993 15 N. J. Turner, *Nat. Chem. Biol.*, 2009, **5**, 567–573.
- 994 16 J. R. Cherry and A. L. Fidantsef, *Curr. Opin. Biotechnol.*, 2003, **14**, 438–443.
- 995 17 K. Rajewsky, H. Gu, R. Kühn, U. A. Betz, W. Müller, J. Roes and F. Schwenk, *J. Clin. Invest.*,  
996 1996, **98**, 600–603.
- 997 18 R. A. Sheldon and S. van Pelt, *Chem. Soc. Rev.*, 2013, **42**, 6223–6235.
- 998 19 R. Dicosimo, J. McAuliffe, A. J. Poulouse and G. Bohlmann, *Chem. Soc. Rev.*, 2013, **42**, 6437–  
999 6474.
- 1000 20 A. Liese and L. Hilterhaus, *Chem. Soc. Rev.*, 2013, **42**, 6236–6249.
- 1001 21 V. M. Balcão and M. M. D. C. Vila, *Adv. Drug Deliv. Rev.*, 2015, **93**, 25–41.
- 1002 22 R. C. Rodrigues, C. Ortiz, A. Berenguer-Murcia, R. Torres and R. Fernández-Lafuente, *Chem.*

- 1003 *Soc. Rev.*, 2013, **42**, 6290–6307.
- 1004 23 C. Garcia-Galan, Á. Berenguer-Murcia, R. Fernandez-Lafuente and R. C. Rodrigues, *Adv.*  
1005 *Synth. Catal.*, 2011, **353**, 2885–2904.
- 1006 24 R. Fernandez-Lafuente, *Enzyme Microb. Technol.*, 2009, **45**, 405–418.
- 1007 25 D. Brady and J. Jordaan, *Biotechnol. Lett.*, 2009, **31**, 1639–1650.
- 1008 26 C. Mateo, J. M. Palomo, G. Fernandez-Lorente, J. M. Guisan and R. Fernandez-Lafuente,  
1009 *Enzyme Microb. Technol.*, 2007, **40**, 1451–1463.
- 1010 27 O. Barbosa, C. Ortiz, Á. Berenguer-Murcia, R. Torres, R. C. Rodrigues and R. Fernandez-  
1011 Lafuente, *Biotechnol. Adv.*, 2015, **33**, 435–456.
- 1012 28 J. M. Woodley, in *Abstract book of the 11th International Conference on Protein Stabilisation*,  
1013 Istanbul, 2016.
- 1014 29 T. Jesionowski, J. Zdarta and B. Krajewska, *Adsorption*, 2014, **20**, 801–821.
- 1015 30 J. C. S. dos Santos, O. Barbosa, C. Ortiz, A. Berenguer-Murcia, R. C. Rodrigues and R.  
1016 Fernandez-Lafuente, *ChemCatChem*, 2015, **7**, 2413–2432.
- 1017 31 E. A. Manoel, J. C. S. dos Santos, D. M. G. Freire, N. Rueda and R. Fernandez-Lafuente,  
1018 *Enzyme Microb. Technol.*, 2015, **71**, 53–57.
- 1019 32 J. C. S. dos Santos, N. Rueda, A. Sanchez, R. Villalonga, L. R. B. Gonçalves and R. Fernandez-  
1020 Lafuente, *RSC Adv.*, 2015, **5**, 35801–35810.
- 1021 33 J. C. S. dos Santos, N. Rueda, R. Torres, O. Barbosa, L. R. B. Gonçalves and R. Fernandez-  
1022 Lafuente, *Process Biochem.*, 2015, **50**, 918–927.
- 1023 34 K.-E. Jaeger, S. Ransac, H. B. Koch, F. Ferrato and B. W. Dijkstra, *FEBS Lett.*, 1993, **332**, 143–  
1024 149.
- 1025 35 J. J. Virgen-Ortíz, V. G. Tacias-Pascacio, D. B. Hirata, B. Torrestiana-Sanchez, A. Rosales-  
1026 Quintero and R. Fernandez-Lafuente, *Enzyme Microb. Technol.*, 2017, **96**, 30–35.
- 1027 36 D. B. Hirata, T. L. Albuquerque, N. Rueda, J. J. Virgen-Ortíz, V. G. Tacias-Pascacio and R.  
1028 Fernandez-Lafuente, *J. Mol. Catal. B Enzym.*, 2016, **133**, 117–123.
- 1029 37 A. Díaz-Rodríguez and B. G. Davis, *Curr. Opin. Chem. Biol.*, 2011, **15**, 211–219.
- 1030 38 B. G. Davis, *Curr. Opin. Biotechnol.*, 2003, **14**, 379–386.
- 1031 39 C. P. Govardhan, *Curr. Opin. Biotechnol.*, 1999, **10**, 331–335.
- 1032 40 S. S. Wong and L. J. C. Wong, *Enzyme Microb. Technol.*, 1992, **14**, 866–874.
- 1033 41 R. Fernandez-Lafuente, C. M. Rosell, G. Alvaro and J. M. Guisan, *Enzyme Microb. Technol.*,  
1034 1992, **14**, 489–495.
- 1035 42 Z. Cabrera, G. Fernandez-Lorente, R. Fernandez-Lafuente, J. M. Palomo and J. M. Guisan,  
1036 *Process Biochem.*, 2009, **44**, 226–231.

- 1037 43 O. Barbosa, M. Ruiz, C. Ortiz, M. Fernández, R. Torres and R. Fernandez-Lafuente, *Process*  
1038 *Biochem.*, 2012, **47**, 867–876.
- 1039 44 C. Garcia-Galan, J. C. S. dos Santos, O. Barbosa, R. Torres, E. B. Pereira, V. C. Corberan, L. R.  
1040 B. Gonçalves and R. Fernandez-Lafuente, *Process Biochem.*, 2014, **49**, 604–616.
- 1041 45 N. Rueda, J. C. S. dos Santos, C. Ortiz, R. Torres, O. Barbosa, R. C. Rodrigues, Á. Berenguer-  
1042 Murcia and R. Fernandez-Lafuente, *Chem. Rec.*, 2016, **16**, 1436–1455.
- 1043 46 R. C. Rodrigues, Á. Berenguer-Murcia and R. Fernandez-Lafuente, *Adv. Synth. Catal.*, 2011,  
1044 **353**, 2216–2238.
- 1045 47 R. C. Rodrigues, C. A. Godoy, G. Volpato, M. A. Z. Ayub, R. Fernandez-Lafuente and J. M.  
1046 Guisan, *Process Biochem.*, 2009, **44**, 963–968.
- 1047 48 B. Brissault, A. Kichler, C. Guis, C. Leborgne, O. Danos and H. Cheradame, *Bioconjug. Chem.*,  
1048 in press, DOI:10.1021/bc0200529.
- 1049 49 S. K. Samal, M. Dash, S. Van Vlierberghe, D. L. Kaplan, E. Chiellini, C. van Blitterswijk, L.  
1050 Moroni and P. Dubruel, *Chem. Soc. Rev.*, 2012, **41**, 7147–7194.
- 1051 50 Q. Yang and T. Runge, *ACS Sustain. Chem. Eng.*, 2016, **4**, 6951–6961.
- 1052 51 S. Hammache, J. S. Hoffman, M. L. Gray, D. J. Fauth, B. H. Howard and H. W. Pennline,  
1053 *Energy Fuels*, 2013, **27**, 6899–6905.
- 1054 52 L. Y. Yurlova, A. P. Kryvoruchko and B. P. Yatsik, *J. Water Chem. Technol.*, 2014, **36**, 115–  
1055 119.
- 1056 53 W. Zhang, H. Liu, C. Sun, T. C. Drage and C. E. Snape, *Chem. Eng. Sci.*, in press,  
1057 DOI:10.1016/j.ces.2014.05.018.
- 1058 54 E. P. Dillon, C. A. Crouse and A. R. Barron, *ACS Nano*, 2008, **2**, 156–164.
- 1059 55 W. Y. Seow, K. Liang, M. Kurisawa and C. A. E. Hauser, *Biomacromolecules*, 2013, **14**, 2340–  
1060 2346.
- 1061 56 S. Sajeesh, T. Y. Lee, S. W. Hong, P. Dua, J. Y. Choe, A. Kang, W. S. Yun, C. Song, S. H.  
1062 Park, S. Kim, C. Li and D. Lee, *Mol. Pharm.*, 2014, **11**, 872–884.
- 1063 57 O. Boussif, F. Lezoualc'h, M. A. Zanta, M. D. Mergny, D. Scherman, B. Demeneix and J. P.  
1064 Behr, *Proc. Natl. Acad. Sci.*, 1995, **92**, 7297–7301.
- 1065 58 A. C. Hunter, *Adv. Drug Deliv. Rev.*, 2006, **58**, 1523–1531.
- 1066 59 C. R. Dick and G. E. Ham, *J. Macromol. Sci. Part A - Chem.*, 1970, **4**, 1301–1314.
- 1067 60 S. Kobayashi, *Prog. Polym. Sci.*, 1990, **15**, 751–823.
- 1068 61 R. Bahulekar, N. R. Ayyangar and S. Ponrathnam, *Enzyme Microb. Technol.*, 1991, **13**, 858–  
1069 868.
- 1070 62 A. Kumar, I. Y. Galaev and B. Mattiasson, *J. Chromatogr. B. Biomed. Sci. Appl.*, 2000, **741**,

- 1071 103–113.
- 1072 63 R. C. Rodrigues, O. Barbosa, C. Ortiz, Á. Berenguer-Murcia, R. Torres and R. Fernandez-  
1073 Lafuente, *RSC Adv.*, 2014, **4**, 38350–38374.
- 1074 64 R. Torres, B. C. C. Pessela, M. Fuentes, C. Mateo, R. Munilla, R. Fernandez-Lafuente and J. M.  
1075 Guisán, *Enzyme Microb. Technol.*, 2006, **39**, 711–716.
- 1076 65 C. Mateo, O. Abian, R. Fernandez-Lafuente and J. M. Guisan, *Biotechnol. Bioeng.*, 2000, **68**,  
1077 98–105.
- 1078 66 M. Fuentes, B. C. C. Pessela, J. V Maquiese, C. Ortiz, R. L. Segura, J. M. Palomo, O. Abian, R.  
1079 Torres, C. Mateo, R. Fernández-Lafuente and J. M. Guisán, *Biotechnol. Prog.*, 2004, **20**, 1134–  
1080 1139.
- 1081 67 B. C. C. Pessela, R. Munilla, L. Betancor, M. Fuentes, A. V Carrascosa, A. Vian, R. Fernandez-  
1082 Lafuente and J. M. Guisán, *J. Chromatogr. A*, 2004, **1034**, 155–159.
- 1083 68 B. C. C. Pessela, L. Betancor, F. Lopez-Gallego, R. Torres, G. M. Dellamora-Ortiz, N. Alonso-  
1084 Morales, M. Fuentes, R. Fernández-Lafuente, J. M. Guisán and C. Mateo, *Enzyme Microb.*  
1085 *Technol.*, 2005, **37**, 295–299.
- 1086 69 A. M. Klibanov, N. O. Kaplan and M. D. Kamen, *Proc. Natl. Acad. Sci. U. S. A.*, 1978, **75**,  
1087 3640–3643.
- 1088 70 B. P. Wasserman, B. S. Jacobson and H. O. Hultin, *Biochim. Biophys. Acta - Enzymol.*, 1981,  
1089 **657**, 52–57.
- 1090 71 L. Gong, H. Dai, S. Zhang and Y. Lin, *Anal. Chem.*, 2016, **88**, 5775–5782.
- 1091 72 L. Li, X. Liu, Q. Xiao, Y. Zheng, S. Wang and B. Feng, *Sens. Lett.*, 2016, **14**, 623–629.
- 1092 73 S. Hosseini, F. Ibrahim, I. Djordjevic, H. A. Rothan, R. Yusof, C. van der Marel and L. H.  
1093 Koole, *Appl. Surf. Sci.*, 2014, **317**, 630–638.
- 1094 74 Y. Yuan, H. He and L. J. Lee, *Biotechnol. Bioeng.*, 2009, **102**, 891–901.
- 1095 75 J. Yakovleva, R. Davidsson, M. Bengtsson, T. Laurell and J. Emnéus, *Biosens. Bioelectron.*,  
1096 2003, **19**, 21–34.
- 1097 76 D. Ivnitski, T. Wolf, B. Solomon, G. Fleminger and J. Rishpon, *Bioelectrochemistry Bioenerg.*,  
1098 1998, **45**, 27–32.
- 1099 77 J. Yakovleva, R. Davidsson, A. Lobanova, M. Bengtsson, S. Eremin, T. Laurell and J. Emnéus,  
1100 *Anal. Chem.*, 2002, **74**, 2994–3004.
- 1101 78 P. Batalla, M. Fuentes, C. Mateo, V. Grazu, R. Fernandez-Lafuente and J. M. Guisan,  
1102 *Biomacromolecules*, 2008, **9**, 2230–2236.
- 1103 79 P. Batalla, M. Fuentes, V. Grazu, C. Mateo, R. Fernandez-Lafuente and J. M. Guisan,  
1104 *Biomacromolecules*, 2008, **9**, 719–723.
- 1105 80 M. Fuentes, C. Mateo, J. M. Guisán and R. Fernández-Lafuente, *Biosens. Bioelectron.*, 2005,



- 1106 20, 1380–1387.
- 1107 81 G. P. Royer, G. M. Green and B. K. Slnha, *J. Macromol. Sci. Part A - Chem.*, 1976, **10**, 289–  
1108 307.
- 1109 82 J. H. T. Luong and K. N. Male, *Enzyme Microb. Technol.*, 1992, **14**, 125–130.
- 1110 83 B. Li, Y. Chen, Z. Cao, H. Niu, D. Liu, Y. He, X. Chen, J. Wu, J. Xie, W. Zhuang and H. Ying,  
1111 *J. Mol. Catal. B Enzym.*, 2014, **101**, 92–100.
- 1112 84 A. V. Barmin, A. V. Eremenko, Ilya N. Kurochkin and T. A. Moskvitina, *Biotechnol. Bioeng.*,  
1113 1994, **44**, 849–853.
- 1114 85 J. C. Pyun, A. S. Jang and J. S. Park, *Enzyme Microb. Technol.*, 1996, **18**, 41–44.
- 1115 86 B. C. . Pessela, R. Fernández-Lafuente, M. Fuentes, A. Vián, J. L. García, A. V. Carrascosa, C.  
1116 Mateo and J. M. Guisán, *Enzyme Microb. Technol.*, 2003, **32**, 369–374.
- 1117 87 P. Gonzalez, F. Batista-Viera and B. M. Brena, *Int. J. Biotechnol.*, 2004, **6**, 338–345.
- 1118 88 S. Pruksasri and S.-T. Yang, in *AIChE Annual Meeting*, 2005, p. 2219.
- 1119 89 R. E. Hamlin, T. L. Dayton, L. E. Johnson and M. S. Johal, *Langmuir*, 2007, **23**, 4432–4437.
- 1120 90 H. A. Güleç, *Colloids Surf. B. Biointerfaces*, 2013, **104**, 83–90.
- 1121 91 A. Martino, M. Durante, P. G. Pifferi, G. Spagna and G. Bianchi, *Process Biochem.*, 1996, **31**,  
1122 281–285.
- 1123 92 M. F. Vieira, A. M. S. Vieira, G. M. Zanin, P. W. Tardioli, C. Mateo and J. M. Guisán, *J. Mol.*  
1124 *Catal. B Enzym.*, 2011, **69**, 47–53.
- 1125 93 M. Filho, B. C. Pessela, C. Mateo, A. V Carrascosa, R. Fernandez-Lafuente and J. M. Guisán,  
1126 *Process Biochem.*, 2008, **43**, 1142–1146.
- 1127 94 V. M. Benassi, T. M. Da Silva, B. C. Pessela, J. M. Guisan, C. Mateo, M. S. Lima, J. A. Jorge  
1128 and M. de L. T. M. Polizeli, *J. Mol. Catal. B Enzym.*, 2013, **89**, 93–101.
- 1129 95 B. C. Dravis, P. E. Swanson and A. J. Russell, *Biotechnol. Bioeng.*, 2001, **75**, 416–423.
- 1130 96 R. Torres, B. C. C. Pessela, C. Mateo, C. Ortiz, M. Fuentes, J. M. Guisan and R. Fernandez-  
1131 Lafuente, *Biotechnol. Prog.*, 2004, **20**, 1297–1300.
- 1132 97 M. Y. Arica and G. Bayramoğlu, *J. Mol. Catal. B Enzym.*, 2006, **38**, 131–138.
- 1133 98 N. Alonso-Morales, F. López-Gallego, L. Betancor, A. Hidalgo, C. Mateo, R. Fernández-  
1134 Lafuente and J. M. Guisán, *Biotechnol. Prog.*, 2008, **20**, 533–536.
- 1135 99 Y. Wei, H. Luo, Y. Chang, H. Yu and Z. Shen, *Biocatal. Biotransformation*, 2015, **33**, 250–259.
- 1136 100 M. Yakup Arica and G. Bayramoğlu, *J. Mol. Catal. B Enzym.*, 2004, **27**, 255–265.
- 1137 101 D. Norouzian, A. Akbarzadeh, S. Mirdamadi, S. Khetami and A. Farhanghi, *Biotechnology*,  
1138 2007, **6**, 436–439.



- 1139 102 C. P. McMahon, G. Rocchitta, P. A. Serra, S. M. Kirwan, J. P. Lowry and R. D. O'Neill,  
1140 *Analyst*, 2006, **131**, 68–72.
- 1141 103 J. M. Bolivar, L. Wilson, S. A. Ferrarotti, R. Fernandez-Lafuente, J. M. Guisan and C. Mateo,  
1142 *Biomacromolecules*, 2006, **7**, 669–673.
- 1143 104 J. M. Bolivar, L. Wilson, S. A. Ferrarotti, R. Fernandez-Lafuente, J. M. Guisan and C. Mateo,  
1144 *Enzyme Microb. Technol.*, 2007, **40**, 540–546.
- 1145 105 X.-P. Wang, B.-J. Gao, J.-F. Guo and L.-P. Zhang, *Gaodeng Xuexiao Huaxue Xuebao/Chemical*  
1146 *J. Chinese Univ.*, 2006, **27**, 1167–1172.
- 1147 106 X.-P. Wang, B.-J. Gao and L.-P. Zhang, *Zhongbei Daxue Xuebao (Ziran Kexue Ban)/Journal*  
1148 *North Univ. China (Natural Sci. Ed.)*, 2007, **28**, 335–340.
- 1149 107 F. Lopez-Gallego, L. Betancor, A. Hidalgo, G. Dellamora-Ortiz, C. Mateo, R. Fernández-  
1150 Lafuente and J. M. Guisán, *Enzyme Microb. Technol.*, 2007, **40**, 278–284.
- 1151 108 N. A. Pchelintsev and P. A. Millner, *Anal. Chim. Acta*, 2008, **612**, 190–197.
- 1152 109 J. Rocha-Martin, D. E. Vega, Z. Cabrera, J. M. Bolivar, R. Fernandez-Lafuente, J. Berenguer  
1153 and J. M. Guisan, *Process Biochem.*, 2009, **44**, 1004–1012.
- 1154 110 V. Sima, C. Cristea, E. Bodoki, G. Duțu and R. Săndulescu, *Cent. Eur. J. Chem.*, 2010, **8**, 1034–  
1155 1040.
- 1156 111 S. F. D'Souza, J. Kumar, S. K. Jha and B. S. Kubal, *Mater. Sci. Eng. C*, 2013, **33**, 850–854.
- 1157 112 A. Ü. Metin, *Macromol. Res.*, 2013, **21**, 1145–1152.
- 1158 113 R. Wu, B. He, G. Zhao and X. Li, *J. Mol. Catal. B Enzym.*, 2014, **99**, 163–168.
- 1159 114 C. Schulz, R. Ludwig and L. Gorton, *Anal. Chem.*, 2014, **86**, 4256–4263.
- 1160 115 B. Ghosh, R. Vaze, A. Anil and A. Lali, *Rasayan J. Chem.*, 2015, **8**, 186–193.
- 1161 116 Y. A. Ramírez Tapias, C. W. Rivero, F. L. Gallego, J. M. Guisán and J. A. Trelles, *Food Chem.*,  
1162 2016, **208**, 252–257.
- 1163 117 T. Montes, V. Grazú, F. López-Gallego, J. A. Hermoso, J. L. García, I. Manso, B. Galán, R.  
1164 González, R. Fernández-Lafuente and J. M. Guisán, *Appl. Environ. Microbiol.*, 2007, **73**, 312–  
1165 319.
- 1166 118 T. Montes, V. Grazu, F. López-Gallego, J. A. Hermoso, J. M. Guisán and R. Fernández-  
1167 Lafuente, *Biomacromolecules*, 2006, **7**, 3052–3058.
- 1168 119 K. S. Atia, *Radiat. Phys. Chem.*, 2005, **73**, 91–99.
- 1169 120 J. J. Virgen-Ortíz, S. Peirce, V. G. Tacias-Pascacio, V. Cortes-Corberan, A. Marzocchella, M. E.  
1170 Russo and R. Fernandez-Lafuente, *Process Biochem.*, 2016, **51**, 1391–1396.
- 1171 121 J. Virgen-Ortíz, S. Pedrero, L. Fernandez-Lopez, N. Lopez-Carrobles, B. Gorines, C. Otero and  
1172 R. Fernandez-Lafuente, *Molecules*, 2017, **22**, 91.

- 1173 122 L. Betancor, F. López-Gallego, A. Hidalgo, N. Alonso-Morales, G. Dellamora-Ortiz, C. Mateo,  
1174 R. Fernández-Lafuente and J. M. Guisán, *Enzyme Microb. Technol.*, 2006, **39**, 877–882.
- 1175 123 F. López-Gallego, L. Betancor, C. Mateo, A. Hidalgo, N. Alonso-Morales, G. Dellamora-Ortiz,  
1176 J. M. Guisán and R. Fernández-Lafuente, *J. Biotechnol.*, 2005, **119**, 70–75.
- 1177 124 O. Barbosa, C. Ortiz, Á. Berenguer-Murcia, R. Torres, R. C. Rodrigues and R. Fernandez-  
1178 Lafuente, *RSC Adv.*, 2014, **4**, 1583–1600.
- 1179 125 O. Barbosa, R. Torres, C. Ortiz and R. Fernandez-Lafuente, *Process Biochem.*, 2012, **47**, 1220–  
1180 1227.
- 1181 126 R. Fernandez-Lafuente, C. M. Rosell, V. Rodriguez and J. M. Guisan, *Enzyme Microb.*  
1182 *Technol.*, 1995, **17**, 517–523.
- 1183 127 J. M. Palomo, R. L. Segura, G. Fernandez-Lorente, R. Fernandez-Lafuente and J. M. Guisán,  
1184 *Enzyme Microb. Technol.*, 2007, **40**, 704–707.
- 1185 128 O. Barbosa, R. Torres, C. Ortiz and R. Fernandez-Lafuente, *Process Biochem.*, 2012, **47**, 766–  
1186 774.
- 1187 129 B. P. Wasserman, D. Burke and B. S. Jacobson, *Enzyme Microb. Technol.*, 1982, **4**, 107–109.
- 1188 130 K. Sankaran, S. S. Godbole and S. F. D'Souza, *Enzyme Microb. Technol.*, 1989, **11**, 617–619.
- 1189 131 K. Furuhashi, S. Deno and M. Sakamoto, *J. Sericultural Sci. Japan*, 1997, **66**, 11–16.
- 1190 132 S. . D'Souza and S. S. Godbole, *J. Biochem. Biophys. Methods*, 2002, **52**, 59–62.
- 1191 133 B. Lakard, G. Herlem, S. Lakard, A. Antoniou and B. Fahys, *Biosens. Bioelectron.*, 2004, **19**,  
1192 1641–1647.
- 1193 134 J. I. S. Valero and S.-T. Yang, in *AIChE Annual Meeting*, 2005, p. 8953.
- 1194 135 N. J. Matella, K. D. Dolan and Y. S. Lee, *J. Food Sci.*, 2006, **71**, C363–C368.
- 1195 136 R. Das, D. Sen, A. Sarkar and C. Bhattacharjee, *J. Inst. Eng. Chem. Eng. Div.*, 2011, **91**, 53–60.
- 1196 137 M. R. El-Aassar, S. S. Al-Deyab and E.-R. Kenawy, *J. Appl. Polym. Sci.*, 2013, **127**, 1873–  
1197 1884.
- 1198 138 A. Nath, C. Bhattacharjee and R. Chowdhury, *Desalination*, 2013, **316**, 31–41.
- 1199 139 F. López-Gallego, L. Betancor, A. Hidalgo, N. Alonso, G. Fernandez-Lorente, J. M. Guisan and  
1200 R. Fernandez-Lafuente, *Enzyme Microb. Technol.*, 2005, **37**, 750–756.
- 1201 140 C. Shu, J. Cai, L. Huang, X. Zhu and Z. Xu, *J. Mol. Catal. B Enzym.*, 2011, **72**, 139–144.
- 1202 141 E. Ondul, N. Dizge and N. Albayrak, *Colloids Surfaces B Biointerfaces*, 2012, **95**, 109–114.
- 1203 142 X. Feng, D. A. Patterson, M. Balaban and E. A. C. Emanuelsson, *Colloids Surfaces B*  
1204 *Biointerfaces*, 2013, **102**, 526–533.
- 1205 143 G. Nicoletti, E. P. Cipolatti, A. Valério, N. G. Carbonera, N. S. Soares, E. Theilacker, J. L.

- 1206 Ninow and D. de Oliveira, *Bioprocess Biosyst. Eng.*, 2015, **38**, 1739–1748.
- 1207 144 N. Kamath, J. S. Melo and S. F. D'Souza, *Appl. Biochem. Biotechnol.*, 1988, **19**, 251–258.
- 1208 145 S. Rocchietti, D. Ubiali, M. Terreni, A. M. Albertini, R. Fernández-Lafuente, J. M. Guisán and  
1209 M. Pregnolato, *Biomacromolecules*, 2004, **5**, 2195–2200.
- 1210 146 A. Fresco-Taboada, I. Serra, J. Fernández-Lucas, C. Acebal, M. Arroyo, M. Terreni and I. de la  
1211 Mata, *Molecules*, 2014, **19**, 11231–11249.
- 1212 147 A. Fresco-Taboada, I. Serra, M. Arroyo, J. Fernández-Lucas, I. De La Mata and M. Terreni,  
1213 *Catal. Today*, 2016, **259**, 197–204.
- 1214 148 G. Bayramoglu, B. Karagoz, N. Bicak and M. Y. Arica, *Ind. Eng. Chem. Res.*, 2014, **53**, 14263–  
1215 14271.
- 1216 149 K. Watanabe and G. P. Royer, *J. Mol. Catal.*, 1983, **22**, 145–152.
- 1217 150 A. B. Salleh, *Biotechnol. Lett.*, 1982, **4**, 769–774.
- 1218 151 E. G. Kondrat'eva, A. N. Reshetilov and A. I. Iaropolov, *Prikl. Biokhim. Mikrobiol.*, 1997, **33**,  
1219 595–599.
- 1220 152 M. M. D. Maia, E. A. Vasconcelos, P. F. C. M. D. Maia, J. C. Maciel, K. R. R. Cajueiro, M. P.  
1221 C. Silva, E. F. Silva Jr, R. A. F. Dutra, V. N. Freire and J. L. Lima Filho, *Process Biochem.*,  
1222 2007, **42**, 429–433.
- 1223 153 I. Kubo, I. Karube, S. Suzuki, Y. Nambu, T. Endo and M. Okawara, *J. Biotechnol.*, 1986, **4**, 27–  
1224 33.
- 1225 154 J. Emnéus and L. Gorton, *Anal. Chim. Acta*, 1993, **276**, 303–318.
- 1226 155 J. Emnéus and L. Gorton, *Anal. Chim. Acta*, 1993, **276**, 319–328.
- 1227 156 A. Lupu, D. Compagnone and G. Palleschi, *Anal. Chim. Acta*, 2004, **513**, 67–72.
- 1228 157 P. Sarkar, *Microchem. J.*, in press, DOI:10.1016/S0026-265X(00)00005-9.
- 1229 158 S. F. D'Souza and B. S. Kubal, *J. Biochem. Biophys. Methods*, 2002, **51**, 151–159.
- 1230 159 T. N. Yudanov, E. S. Obolonkova, T. A. Cherdyntseva, A. I. Netrusov and L. S. Gal'braikh,  
1231 *Fibre Chem.*, 2004, **36**, 122–125.
- 1232 160 N. Alonso, F. López-Gallego, L. Betancor, A. Hidalgo, C. Mateo, J. M. Guisan and R.  
1233 Fernandez-Lafuente, *J. Mol. Catal. B Enzym.*, 2005, **35**, 57–61.
- 1234 161 B. Gao, X. Wang and Y. Shen, *Biochem. Eng. J.*, 2006, **28**, 140–147.
- 1235 162 A. Caro, V. Humblot, C. Méthivier, M. Minier, L. Barbes, J. Li, M. Salmain and C.-M. Pradier,  
1236 *J. Colloid Interface Sci.*, 2010, **349**, 13–18.
- 1237 163 R. Xu, C. Chi, F. Li and B. Zhang, *Bioresour. Technol.*, 2013, **149**, 111–116.
- 1238 164 G. B. Oliveira, J. L. Lima Filho, M. E. Cavalcante Chaves, W. M. Azevedo and L. B. Carvalho,

- 1239 *React. Funct. Polym.*, 2008, **68**, 27–32.
- 1240 165 P. G. Cadena, R. A. S. Jeronimo, J. M. Melo, R. A. Silva, J. L. Lima Filho and M. C. B.  
1241 Pimentel, *Bioresour. Technol.*, 2010, **101**, 1595–1602.
- 1242 166 M. I. Wahba and M. E. Hassan, *Biopolymers*, 2015, **103**, 675–684.
- 1243 167 C. Cui, Y. Tao, L. Li, B. Chen and T. Tan, *J. Mol. Catal. B Enzym.*, 2013, **91**, 59–66.
- 1244 168 J. D. An, D. A. Patterson, S. McNeil and M. M. Hossain, *Biotechnol. Prog.*, 2014, **30**, 806–817.
- 1245 169 L. Zhao, Q. Liu, S. Yan, Z. Chen, J. Chen and X. Li, *J. Biotechnol.*, 2013, **168**, 46–54.
- 1246 170 A. M. Hashem, A. A. Gamal, M. E. Hassan, N. M. Hassanein and M. A. Esawy, *Int. J. Biol.*  
1247 *Macromol.*, 2016, **82**, 905–912.
- 1248 171 N. C. Dubey, B. P. Tripathi, M. Müller, M. Stamm and L. Ionov, *Biomacromolecules*, 2016, **17**,  
1249 1610–1620.
- 1250 172 S. Ates and S. Pekyardimci, *J. Macromol. Sci. Part A*, 1995, **32**, 337–345.
- 1251 173 S. A. Costa and R. L. Reis, *J. Mater. Sci. Mater. Med.*, 2004, **15**, 335–342.
- 1252 174 B. Akkaya, F. Şahin, G. Demirel and H. Tümtürk, *Biochem. Eng. J.*, 2009, **43**, 333–337.
- 1253 175 M. Khoobi, S. F. Motevalizadeh, Z. Asadgol, H. Forootanfar, A. Shafiee and M. A. Faramarzi,  
1254 *Biochem. Eng. J.*, 2014, **88**, 131–141.
- 1255 176 J. Zemek, L. Kuniak, P. Gemeiner, J. Zámocký and Š. Kučár, *Enzyme Microb. Technol.*, 1982,  
1256 **4**, 233–238.
- 1257 177 Y. Hatanaka, T. Tomohiro, N. Tachi and Y. Azuma, *Heterocycles*, 2009, **79**, 897.
- 1258 178 C. L. Chuang, Y. J. Wang and H. L. Lan, *Anal. Chim. Acta*, 1997, **353**, 37–44.
- 1259 179 R. A. Bohara, N. D. Thorat and S. H. Pawar, *RSC Adv.*, 2016, **6**, 43989–44012.
- 1260 180 E. T. Hwang and M. B. Gu, *Eng. Life Sci.*, 2013, **13**, 49–61.
- 1261 181 S. A. Ansari and Q. Husain, *Biotechnol. Adv.*, 2012, **30**, 512–523.
- 1262 182 J. M. Bolivar, S. Schelch, T. Mayr and B. Nidetzky, *ChemCatChem*, 2014, **6**, 981–986.
- 1263 183 J. M. Bolivar, T. Consolati, T. Mayr and B. Nidetzky, *Biotechnol. Bioeng.*, 2013, **110**, 2086–  
1264 2095.
- 1265 184 C. Boniello, T. Mayr, J. M. Bolivar and B. Nidetzky, *BMC Biotechnol.*, 2012, **12**, article 11.
- 1266 185 J. M. Guisan, G. Alvaro, C. Rosell and R. Fernandez-Lafuente, *Biotechnol. Appl. Biochem.*,  
1267 1994, **20**, 357–369.
- 1268 186 L. Betancor, M. Fuentes, G. Dellamora-Ortiz, F. López-Gallego, A. Hidalgo, N. Alonso-  
1269 Morales, C. Mateo, J. M. Guisán and R. Fernández-Lafuente, *J. Mol. Catal. B Enzym.*, 2005, **32**,  
1270 97–101.

- 1271 187 L. Betancor, F. López-Gallego, A. Hidalgo, N. Alonso-Morales, M. Fuentes, R. Fernández-  
1272 Lafuente and J. M. Guisán, *J. Biotechnol.*, 2004, **110**, 201–207.
- 1273 188 J. M. Bolivar, J. Rocha-Martin, C. Mateo, F. Cava, J. Berenguer, R. Fernandez-Lafuente and J.  
1274 M. Guisan, *Biomacromolecules*, 2009, **10**, 742–747.
- 1275 189 C. Garcia-Galan, O. Barbosa and R. Fernandez-Lafuente, *Enzyme Microb. Technol.*, 2013, **52**,  
1276 211–217.
- 1277 190 E. P. Cipolatti, R. O. Henriques, D. E. Moritz, J. L. Ninow, D. M. G. Freire, E. A. Manoel and  
1278 R. Fernandez-, *RSC Adv.*, 2016, **6**, 104675–104692.
- 1279 191 E. P. Cipolatti, M. J. A. Silva, M. Klein, V. Feddern, M. M. C. Feltes, J. V Oliveira, J. L. Ninow  
1280 and D. De Oliveira, *J. Mol. Catal. B Enzym.*, 2014, **99**, 56–67.
- 1281 192 S. Feijoo, S. González-García, Y. Moldes-Diz, C. Vazquez-Vazquez, G. Feijoo and M. T.  
1282 Moreira, *J. Clean. Prod.*, 2017, **143**, 528–538.
- 1283 193 J. Xu, F. Zeng, S. Wu, X. Liu, C. Hou and Z. Tong, *Nanotechnology*, 2007, **18**, 265704.
- 1284 194 B. Sahoo, S. Dutta and D. Dhara, *J. Chem. Sci.*, 2016, **128**, 1131–1140.
- 1285 195 K. Solanki and M. N. Gupta, *New J. Chem.*, 2011, **35**, 2551–2556.
- 1286 196 J. Mukherjee, K. Solanki and M. N. Gupta, in *Immobilization of Enzymes and Cells*, ed. J. M.  
1287 Guisan, Humana Press, New York, Third Edit., 2013, vol. 1051, pp. 117–127.
- 1288 197 V. Patel, C. Shah, M. Deshpande and D. Madamwar, *Appl. Biochem. Biotechnol.*, 2016, **178**,  
1289 1630–1651.
- 1290 198 M. Khoobi, S. F. Motevalizadeh, Z. Asadgol, H. Forootanfar, A. Shafiee and M. A. Faramarzi,  
1291 *Mater. Chem. Phys.*, 2015, **149**, 77–86.
- 1292 199 S. F. Motevalizadeh, M. Khoobi, A. Sadighi, M. Khalilvand-Sedagheh, M. Pazhouhandeh, A.  
1293 Ramazani, M. A. Faramarzi and A. Shafiee, *J. Mol. Catal. B Enzym.*, 2015, **120**, 75–83.
- 1294 200 H. Shen, J. Hou, Z. Gao, B. Chen and T. Tan, *Appl. Biochem. Biotechnol.*, 2014, **173**, 2279–  
1295 2286.
- 1296 201 M. Khoobi, M. Khalilvand-Sedagheh, A. Ramazani, Z. Asadgol, H. Forootanfar and M. A.  
1297 Faramarzi, *J. Chem. Technol. Biotechnol.*, 2016, **91**, 375–384.
- 1298 202 J.-F. Liu, H. Liu, B. Tan, Y.-H. Chen and R.-J. Yang, *J. Mol. Catal. B Enzym.*, 2012, **82**, 64–70.
- 1299 203 M. Minig, L. S. Mazzaferro, A. Capece and J. D. Breccia, *Biotechnol. Appl. Biochem.*, 2015,  
1300 **62**, 94–100.
- 1301 204 C. Monteil, N. Bar, R. Retoux, J. Henry, B. Bernay and D. Villemin, *Sensors Actuators, B*  
1302 *Chem.*, 2014, **192**, 269–274.
- 1303 205 Y. Zhang, H. Ren, Y. Wang, K. Chen, B. Fang and S. Wang, *Sci. Rep.*, 2016, **6**, 24163.
- 1304 206 T. T. Xia, C. Z. Liu, J. H. Hu and C. Guo, *Chem. Eng. J.*, 2016, **295**, 201–206.

- 1305 207 F. Zhang, R. Wang, C. Zhen and B. Li, *J. Mol. Catal. B Enzym.*, 2016, **134**, 164–171.
- 1306 208 H. Tümtürk and H. Yüksekdağ, *Artif. Cells, Nanomedicine, Biotechnol.*, 2016, **44**, 443–447.
- 1307 209 P. Zamora, A. Narváez and E. Domínguez, *Bioelectrochemistry*, 2009, **76**, 100–106.
- 1308 210 H. Ren, Y. Zhang, J. Su, P. Lin, B. Wang, B. Fang and S. Wang, *J. Biotechnol.*, 2017, **241**, 33–  
1309 41.
- 1310 211 M. Müller and B. Urban, *Macromol. Chem. Phys.*, 2016, **217**, 2360–2371.
- 1311 212 T. Zhang, W. Wang, D. Zhang, X. Zhang, M. Yurong, Y. Zhou and L. Qi, *Adv. Funct. Mater.*,  
1312 2010, **20**, 1152–1160.
- 1313 213 V. Incani, C. Danumah and Y. Boluk, *Cellulose*, 2013, **20**, 191–200.
- 1314 214 S. Zhao, X. Ji, P. Lin and Y. M. Liu, *Anal. Biochem.*, 2011, **411**, 88–93.
- 1315 215 A. Gutiérrez, M. C. Rodríguez, L. Galicia and G. A. Rivas, *ECS Trans.*, 2011, **36**, 393–400.
- 1316 216 S. Zhang, Z. Jiang, W. Zhang, X. Wang and J. Shi, *Biochem. Eng. J.*, 2015, **93**, 281–288.
- 1317 217 N. Deng, B. Jiang, Y. Chen, Z. Liang, L. Zhang, Y. Liang, K. Yang and Y. Zhang, *J.*  
1318 *Chromatogr. A*, 2016, **1427**, 16–21.
- 1319 218 Y. M. Yan, I. Baravik, O. Yehezkeli and I. Willner, *J. Phys. Chem. C*, 2008, **112**, 17883–17888.
- 1320 219 D. Brondani, B. de Souza, B. S. Souza, A. Neves and I. C. Vieira, *Biosens. Bioelectron.*, 2013,  
1321 **42**, 242–247.
- 1322 220 O. A. Loaiza, P. J. Lamas-Ardisana, L. Añorga, E. Jubete, V. Ruiz, M. Borghei, G. Cabañero  
1323 and H. J. Grande, *Bioelectrochemistry*, 2015, **101**, 58–65.
- 1324 221 S. F. D'Souza and J. S. Melo, *Enzyme Microb. Technol.*, 1991, **13**, 508–511.
- 1325 222 Y. Bai and S. T. Yang, *Biotechnol. Bioeng.*, 2005, **92**, 137–146.
- 1326 223 T. Zhu and K. H. Row, *J. Sep. Sci.*, 2012, **35**, 1294–1302.
- 1327 224 A. Ghanem and T. Ikegami, *J. Sep. Sci.*, 2011, **34**, 1945–1957.
- 1328 225 C. Delattre and M. A. Vijayalakshmi, *J. Mol. Catal. B Enzym.*, 2009, **60**, 97–105.
- 1329 226 J. Krenkova and F. Svec, *J. Sep. Sci.*, 2009, **32**, 706–718.
- 1330 227 J. Ma, L. Zhang, Z. Liang, W. Zhang and Y. Zhang, *J. Sep. Sci.*, 2007, **30**, 3050–3059.
- 1331 228 F. Svec, *Electrophoresis*, 2006, **27**, 947–961.
- 1332 229 A. Jungbauer and R. Hahn, *J. Sep. Sci.*, 2004, **27**, 767–778.
- 1333 230 Y. Zheng, X. Wang and Y. Ji, *Talanta*, 2012, **91**, 7–17.
- 1334 231 R. Mallik and D. S. Hage, *J. Sep. Sci.*, 2006, **29**, 1686–1704.



- 1335 232 S. Xie, R. W. Allington, J. M. J. Fréchet and F. Svec, *Adv. Biochem. Eng. Biotechnol.*, 2002, **76**,  
1336 87–125.
- 1337 233 K. M. De Lathouder, J. Bakker, M. T. Kreutzer, F. Kapteijn, J. A. Moulijn and S. A. Wallin,  
1338 *Chem. Eng. Sci.*, 2004, **59**, 5027–5033.
- 1339 234 K. M. de Lathouder, D. T. J. van Benthem, S. A. Wallin, C. Mateo, R. F. Lafuente, J. M.  
1340 Guisan, F. Kapteijn and J. A. Moulijn, *J. Mol. Catal. B Enzym.*, 2008, **50**, 20–27.
- 1341 235 Y. Q. Lv, D. Y. Fu, T. W. Tan and M. Y. Wang, *J. Mol. Catal. B Enzym.*, 2010, **62**, 149–154.
- 1342 236 P. He, J. Davies, G. Greenway and S. J. Haswell, *Anal. Chim. Acta*, 2010, **659**, 9–14.
- 1343 237 O. Zaushitsyna, D. Berillo, H. Kirsebom and B. Mattiasson, *Top. Catal.*, 2014, **57**, 339–348.
- 1344 238 D. Yang, J. Zhao, X. Wang, J. Shi, S. Zhang and Z. Jiang, *Biochem. Eng. J.*, 2017, **117**, 52–61.
- 1345 239 P. He, G. Greenway and S. J. Haswell, *Microfluid. Nanofluidics*, 2010, **8**, 565–573.
- 1346 240 D. Ogónczyk, P. Jankowski and P. Garstecki, *Lab Chip*, 2012, **12**, 2743–2748.
- 1347 241 L. M. Cerdeira Ferreira, E. T. da Costa, C. L. do Lago and L. Angnes, *Biosens. Bioelectron.*,  
1348 2013, **47**, 539–544.
- 1349 242 M. R. F. Cerqueira, D. Grasseschi, R. C. Matos and L. Angnes, *Talanta*, 2014, **126**, 20–26.
- 1350 243 M. R. F. Cerqueira, M. S. F. Santos, R. C. Matos, I. G. R. Gutz and L. Angnes, *Microchem. J.*,  
1351 2015, **118**, 231–237.
- 1352 244 P. Lozano, A. B. Pérez-Marín, T. De Diego, D. Gómez, D. Paolucci-Jeanjean, M. P. Belleville,  
1353 G. M. Rios and J. L. Iborra, *J. Memb. Sci.*, 2002, **201**, 55–64.
- 1354 245 Y. Liu, H. Wang, J. Huang, J. Yang, B. Liu and P. Yang, *Anal. Chim. Acta*, 2009, **650**, 77–82.
- 1355 246 G. Roman-Gusetu, K. C. Waldron and D. Rochefort, *J. Chromatogr. A*, 2009, **1216**, 8270–8276.
- 1356 247 N. G. Patel, A. Erlenkötter, K. Cammann and G. C. Chemnitz, *Sensors Actuators, B Chem.*,  
1357 2000, **67**, 134–141.
- 1358 248 M. Onda, K. Ariga and T. Kunitake, *J. Biosci. Bioeng.*, 1999, **87**, 69–75.
- 1359 249 M. C. Rodríguez and G. A. Rivas, *Electroanalysis*, 2004, **16**, 1717–1722.
- 1360 250 C. Deng, J. Chen, Z. Nie and S. Si, *Biosens. Bioelectron.*, 2010, **26**, 213–219.
- 1361 251 N. Albayrak and S.-T. T. Yang, *Biotechnol. Prog.*, 2002, **18**, 240–251.
- 1362 252 H. A. Güleç, S. Gürdaş, N. Albayrak and M. Mutlu, *Biotechnol. Bioprocess Eng.*, 2010, **15**,  
1363 1006–1015.
- 1364 253 D. E. Wong, J. N. Talbert and J. M. Goddard, *J. Food Sci.*, 2013, **78**, E853–E860.
- 1365 254 S. R. Forrest, B. B. Elmore and J. D. Palmer, *Appl. Biochem. Biotechnol.*, 2005, **121**, 85–91.
- 1366 255 S. R. Forrest, B. B. Elmore and J. D. Palmer, *Catal. Today*, 2007, **120**, 30–34.

- 1367 256 F. Bucatariu, C. A. Ghiorghita, F. Simon, C. Bellmann and E. S. Dragan, *Appl. Surf. Sci.*, 2013,  
1368 **280**, 812–819.
- 1369 257 J. J. Karimpil, J. S. Melo and S. F. D'Souza, *Int. J. Biol. Macromol.*, 2012, **50**, 300–302.
- 1370 258 J. M. Palomo, M. Fuentes, G. Fernández-Lorente, C. Mateo, J. M. Guisan and R. Fernández-  
1371 Lafuente, *Biomacromolecules*, 2003, **4**, 1–6.
- 1372 259 J. M. Palomo, C. Ortiz, G. Fernández-Lorente, M. Fuentes, J. M. Guisán and R. Fernández-  
1373 Lafuente, *Enzyme Microb. Technol.*, 2005, **36**, 447–454.
- 1374 260 Y. Lvov, R. Price, B. Gaber and I. Ichinose, *Colloids Surfaces A Physicochem. Eng. Asp.*, 2002,  
1375 **198–200**, 375–382.
- 1376 261 M. Onda, Y. Lvov, K. Ariga and T. Kunitake, *J. Ferment. Bioeng.*, 1996, **82**, 502–506.
- 1377 262 J. P. Santos, E. R. Welsh, B. P. Gaber and A. Singh, *Langmuir*, 2001, **17**, 5361–5367.
- 1378 263 M. K. Ram, P. Bertoncello, H. Ding, S. Paddeu and C. Nicolini, *Biosens. Bioelectron.*, 2001, **16**,  
1379 849–856.
- 1380 264 A. Uzunoglu, I. Ramirez, E. Andreasen and L. A. Stanciu, *Microchim. Acta*, 2016, **183**, 1667–  
1381 1675.
- 1382 265 S. Peirce, J. J. Virgen-Ortiz, V. G. Tacias-Pascacio, N. Rueda, R. Bartolome-Cabrero, L.  
1383 Fernandez-Lopez, M. E. Russo, A. Marzocchella and R. Fernandez-Lafuente, *RSC Adv.*, 2016,  
1384 **6**, 61707–61715.
- 1385 266 H. Zaak, L. Fernandez-Lopez, V. Cortes-Corberan, J. F. Kornecki, E.-H. Siar and R. Fernandez-  
1386 Lafuente, in press, DOI 10.1016/j.procbio.2017.06.014.
- 1387 267 M. Kierstan and C. Bucke, *Biotechnol. Bioeng.*, 1977, **19**, 387–397.
- 1388 268 E. Taqieddin and M. Amiji, *Biomaterials*, 2004, **25**, 1937–1945.
- 1389 269 C.-H. Fan and C.-K. Lee, *Biochem. Eng. J.*, 2001, **8**, 157–164.
- 1390 270 M. M. M. Elnashar, M. A. Yassin and T. Kahil, *J. Appl. Polym. Sci.*, 2008, **109**, 4105–4111.
- 1391 271 A. C. Pierre, *Biocatal. Biotransformation*, 2004, **22**, 145–170.
- 1392 272 H. Yang, S. G. Cao, L. Ma, Z. T. Ding, S. D. Liu and Y. H. Cheng, *Biochem. Biophys. Res.*  
1393 *Commun.*, 1994, **200**, 83–88.
- 1394 273 P. D. Desai, A. M. Dave and S. Devi, *J. Mol. Catal. B Enzym.*, 2004, **31**, 143–150.
- 1395 274 I. S. Tan and K. T. Lee, *Bioresour. Technol.*, 2015, **184**, 386–394.
- 1396 275 M. I. Wahba, *Int. J. Biol. Macromol.*, 2016, **91**, 877–886.
- 1397 276 C. Chang, D. E. Ward, B. Kelemen and J. C. McAuliffe, in *Biomolecular Catalysis*, eds. J. Kim,  
1398 S. H. Kim and P. Wang, American Chemical Society, Washington, DC, 2008, vol. 986, pp. 183–  
1399 198.



- 1400 277 N. A. Pchelintsev, F. Neville and P. A. Millner, *Sensors Actuators B Chem.*, 2008, **135**, 21–26.
- 1401 278 N. C. Dubey, B. P. Tripathi, M. Müller, M. Stamm and L. Ionov, *ACS Appl. Mater. Interfaces*,  
1402 2015, **7**, 1500–1507.
- 1403 279 S. Zhang, Z. Jiang, X. Wang, C. Yang and J. Shi, *ACS Appl. Mater. Interfaces*, 2015, **7**, 19570–  
1404 19578.
- 1405 280 J. M. Qian, A. L. Suo, Y. Yao and Z. H. Jin, *Clin. Biochem.*, 2004, **37**, 155–161.
- 1406 281 D. Rochefort, L. Kouisni and K. Gendron, *J. Electroanal. Chem.*, 2008, **617**, 53–63.
- 1407 282 M. Hébert and D. Rochefort, *ECS Trans.*, 2008, **16**, 85–97.
- 1408 283 Y. Zhang and D. Rochefort, *J. Microencapsul.*, 2010, **27**, 703–713.
- 1409 284 Y. Zhang and D. Rochefort, *Process Biochem.*, 2011, **46**, 993–1000.
- 1410 285 A. Savolainen, Y. Zhang, D. Rochefort, U. Holopainen, T. Erho, J. Virtanen and M. Smolander,  
1411 *Biomacromolecules*, 2011, **12**, 2008–2015.
- 1412 286 R. Huang, M. Wu, M. J. Goldman and Z. Li, *Biotechnol. Bioeng.*, 2015, **112**, 1092–1101.
- 1413 287 L. Cao, F. Van Rantwijk and R. A. Sheldon, *Org. Lett.*, 2000, **2**, 1361–1364.
- 1414 288 R. Schoevaart, M. W. Wolbers, M. Golubovic, M. Ottens, A. P. G. Kieboom, F. Van Rantwijk,  
1415 L. A. M. Van Der Wielen and R. A. Sheldon, *Biotechnol. Bioeng.*, 2004, **87**, 754–762.
- 1416 289 D. Häring and P. Schreier, *Curr. Opin. Chem. Biol.*, 1999, **3**, 35–38.
- 1417 290 E. K. Yan, H. L. Cao, C. Y. Zhang, Q. Q. Lu, Y. J. Ye, J. He, L. J. Huang and D. C. Yin, *RSC*  
1418 *Adv.*, 2015, **5**, 26163–26174.
- 1419 291 R. A. Sheldon, *Appl. Microbiol. Biotechnol.*, 2011, **92**, 467–477.
- 1420 292 R. A. Sheldon, *Org. Process Res. Dev.*, 2011, **15**, 213–223.
- 1421 293 A. Chmura, S. Rustler, M. Paravidino, F. Van Rantwijk, A. Stolz and R. A. Sheldon,  
1422 *Tetrahedron Asymmetry*, 2013, **24**, 1225–1232.
- 1423 294 L. Dal Magro, P. F. Hertz, R. Fernandez-Lafuente, M. P. Klein and R. C. Rodrigues, *RSC Adv.*,  
1424 2016, **6**, 27242–27251.
- 1425 295 K. Ahumada, A. Martínez-Gil, Y. Moreno-Simunovic, A. Illanes and L. Wilson, *Molecules*,  
1426 2016, **21**, 1485.
- 1427 296 Q. Zhang, X. Zha, N. Zhou and Y. Tian, *J. Basic Microbiol.*, 2016, **56**, 422–431.
- 1428 297 J. D. Cui, R. L. Liu and L. B. Li, *Korean J. Chem. Eng.*, 2016, **33**, 610–615.
- 1429 298 J. Cruz, O. Barbosa, R. C. Rodrigues, R. Fernandez-Lafuente, R. Torres and C. Ortiz, *J. Mol.*  
1430 *Catal. B Enzym.*, 2012, **80**, 7–14.
- 1431 299 M. Galvis, O. Barbosa, M. Ruiz, J. Cruz, C. Ortiz, R. Torres and R. Fernandez-Lafuente,

- 1432 *Process Biochem.*, 2012, **47**, 2373–2378.
- 1433 300 A. C. O. Mafra, W. Kopp, M. B. Beltrame, R. de Lima Camargo Giordano, M. P. de Arruda  
1434 Ribeiro and P. W. Tardioli, *J. Mol. Catal. B Enzym.*, 2016, **133**, 107–116.
- 1435 301 F. López-Gallego, L. Betancor, A. Hidalgo, N. Alonso, R. Fernández-Lafuente and J. M.  
1436 Guisán, *Biomacromolecules*, 2005, **6**, 1839–1842.
- 1437 302 L. Wilson, G. Fernández-Lorente, R. Fernández-Lafuente, A. Illanes, J. M. Guisán and J. M.  
1438 Palomo, *Enzyme Microb. Technol.*, 2006, **39**, 750–755.
- 1439 303 J. Yan, X. Gui, G. Wang and Y. Yan, *Appl. Biochem. Biotechnol.*, 2012, **166**, 925–932.
- 1440 304 J. Pan, X. D. Kong, C. X. Li, Q. Ye, J. H. Xu and T. Imanaka, *J. Mol. Catal. B Enzym.*, 2011,  
1441 **68**, 256–261.
- 1442 305 C. Mateo, B. Fernandes, F. Van Rantwijk, A. Stolz and R. A. Sheldon, *J. Mol. Catal. B Enzym.*,  
1443 2006, **38**, 154–157.
- 1444 306 L. Wilson, A. Illanes, O. Abián, B. C. C. Pessela, R. Fernández-Lafuente and J. M. Guisán,  
1445 *Biomacromolecules*, 2004, **5**, 852–857.
- 1446 307 O. Romero, J. Vergara, R. Fernández-Lafuente, J. M. Guisán, A. Illanes and L. Wilson,  
1447 *Biotechnol. Bioeng.*, 2009, **103**, 472–479.
- 1448 308 L. Wilson, A. Illanes, L. Soler and M. J. Henríquez, *Process Biochem.*, 2009, **44**, 322–326.
- 1449 309 L. Wilson, A. Illanes, O. Romero, J. Vergara and C. Mateo, *Enzyme Microb. Technol.*, 2008, **43**,  
1450 442–447.
- 1451 310 O. M. Poltorak, E. S. Chukhray and I. Y. Torshin, *Biochem.*, 1998, **63**, 303–311.
- 1452 311 R. W. Lencki, J. Arul and R. J. Neufeld, *Biotechnol. Bioeng.*, 1992, **40**, 1427–1434.
- 1453 312 O. M. Poltorak, E. S. Chukhrai and I. Y. Torshin, *Russ. J. Phys. Chem. A*, 2000, **74**, S400–S410.
- 1454 313 R. Fernández-Lafuente, V. Rodríguez, C. Mateo, G. Penzol, O. Hernández-Justiz, G. Irazoqui,  
1455 A. Villarino, K. Ovsejevi, F. Batista and J. M. Guisán, *J. Mol. Catal. B Enzym.*, 1999, **7**, 181–  
1456 189.
- 1457 314 M. Fuentes, R. L. Segura, O. Abian, L. Betancor, A. Hidalgo, C. Mateo, R. Fernandez-Lafuente  
1458 and J. M. Guisan, *Proteomics*, 2004, **4**, 2602–2607.
- 1459 315 I. Serra, T. Bavaro, D. A. Cecchini, S. Daly, A. M. Albertini, M. Terreni and D. Ubiali, *J. Mol.*  
1460 *Catal. B Enzym.*, 2013, **95**, 16–22.
- 1461 316 I. Serra, D. Ubiali, J. Piškur, S. Christoffersen, E. S. Lewkowicz, A. M. Iribarren, A. M.  
1462 Albertini and M. Terreni, *Chempluschem*, 2013, **78**, 157–165.
- 1463 317 I. Serra, C. D. Serra, S. Rocchietti, D. Ubiali and M. Terreni, *Enzyme Microb. Technol.*, 2011,  
1464 **49**, 52–58.
- 1465 318 D. Ubiali, S. Rocchietti, F. Scaramozzino, M. Terreni, A. M. Albertini, R. Fernández-Lafuente,

- 1466 J. M. Guisán and M. Pregmolato, *Adv. Synth. Catal.*, 2004, **346**, 1361–1366.
- 1467 319 C. R. F. Terrasan, C. C. Aragon, D. C. Masui, B. C. Pessela, G. Fernandez-Lorente, E. C.  
1468 Carmona and J. M. Guisan, *Biocatal. Biotransformation*, 2016, **34**, 161–171.
- 1469 320 J. M. Bolivar, J. Rocha-Martín, C. Mateo and J. M. Guisan, *Process Biochem.*, 2012, **47**, 679–  
1470 686.
- 1471 321 H. He, Y. Wei, H. Luo, X. Li, X. Wang, C. Liang, Y. Chang, H. Yu and Z. Shen, *Biotechnol.*  
1472 *Prog.*, 2015, **31**, 387–395.
- 1473 322 O. Abian, L. Wilson, C. Mateo, G. Fernández-Lorente, J. M. Palomo, R. Fernández-Lafuente, J.  
1474 M. Guisán, D. Re, A. Tam and M. Daminatti, *J. Mol. Catal. B Enzym.*, 2002, **19–20**, 295–303.
- 1475 323 O. Abian, C. Mateo, G. Fernández-Lorente, J. M. Palomo, R. Fernández-Lafuente and J. M.  
1476 Guisán, *Biocatal. Biotransformation*, 2001, **19**, 489–503.
- 1477 324 O. Abian, C. Mateo, G. Fernández-Lorente, J. M. Guisán and R. Fernández-Lafuente,  
1478 *Biotechnol. Prog.*, 2003, **19**, 1639–1642.
- 1479 325 G. Volpato, R. C. Rodrigues and R. Fernandez-Lafuente, *Curr. Med. Chem.*, 2010, **17**, 3855–  
1480 3873.
- 1481 326 O. Abian, C. Mateo, G. Fernández-Lorente, J. M. Guisán and R. Fernández-Lafuente,  
1482 *Biotechnol. Prog.*, 2004, **20**, 117–121.
- 1483 327 O. Abian, C. Mateo, J. M. Palomo, G. Fernández-Lorente, J. M. Guisán and R. Fernández-  
1484 Lafuente, *Biotechnol. Prog.*, 2004, **20**, 984–988.
- 1485 328 M. Fuentes, P. Batalla, V. Grazu, B. C. C. Pessela, C. Mateo, T. Montes, J. A. Hermoso, J. M.  
1486 Guisan and R. Fernandez-Lafuente, *Biomacromolecules*, 2007, **8**, 703–707.
- 1487 329 T. Montes, V. Grazú, I. Manso, B. Galán, F. López-Gallego, R. González, J. A. Hermoso, J. L.  
1488 García, J. M. Guisán and R. Fernández-Lafuente, *Adv. Synth. Catal.*, 2007, **349**, 459–464.
- 1489 330 J. M. Guisan, P. Sabuquillo, R. Fernandez-Lafuente, G. Fernandez-Lorente, C. Mateo, P. J.  
1490 Halling, D. Kennedy, E. Miyata and D. Re, *J. Mol. Catal. B Enzym.*, 2001, **11**, 817–824.
- 1491 331 M. Villalba, C. M. Verdasco-Martín, J. C. S. dos Santos, R. Fernandez-Lafuente and C. Otero,  
1492 *Enzyme Microb. Technol.*, 2016, **90**, 35–44.
- 1493 332 S. Moreno-Perez, M. Filice, J. M. Guisan and G. Fernandez-Lorente, *Chem. Phys. Lipids*, 2013,  
1494 **174**, 48–54.
- 1495 333 J. M. Palomo, G. Muoz, G. Fernández-Lorente, C. Mateo, R. Fernández-Lafuente and J. M.  
1496 Guisán, *J. Mol. Catal. B Enzym.*, 2002, **19–20**, 279–286.
- 1497 334 K. K. Kim, H. K. Song, D. H. Shin, K. Y. Hwang and S. W. Suh, *Structure*, 1997, **5**, 173–185.
- 1498 335 M. Cygler and J. D. Schrag, *Biochim. Biophys. Acta*, 1999, **1441**, 205–214.
- 1499 336 N. Rueda, J. C. S. dos Santos, R. Torres, C. Ortiz, O. Barbosa and R. Fernandez-Lafuente, *RSC*  
1500 *Adv.*, 2015, **5**, 11212–11222.

- 1501 337 D. B. Hirata, T. L. Albuquerque, N. Rueda, J. M. Sánchez-Montero, E. Garcia-Verdugo, R.  
1502 Porcar and R. Fernandez-Lafuente, *ChemistrySelect*, 2016, **1**, 3259–3270.
- 1503 338 C. Bernal, A. Illanes and L. Wilson, *Langmuir*, 2014, **30**, 3557–3566.
- 1504 339 C. Bernal, A. Illanes and L. Wilson, *J. Agric. Food Chem.*, 2015, **63**, 3716–3724.
- 1505 340 N. Guajardo, C. Bernal, L. Wilson and Z. Cabrera, *Catal. Today*, 2015, **255**, 21–26.
- 1506 341 N. Guajardo, C. Bernal, L. Wilson and Z. Cabrera, *Process Biochem.*, 2015, **50**, 1870–1877.
- 1507 342 V. Vescovi, R. Giordano, A. Mendes and P. Tardioli, *Molecules*, 2017, **22**, 212.
- 1508 343 G. Fernandez-Lorente, M. Filice, D. Lopez-Vela, C. Pizarro, L. Wilson, L. Betancor, Y. Avila  
1509 and J. M. Guisan, *J. Am. Oil Chem. Soc.*, 2011, **88**, 801–807.
- 1510 344 L. Fernandez-Lopez, S. G. Pedrero, N. Lopez-Carrobles, J. J. Virgen-Ortíz, B. C. Gorines, C.  
1511 Otero and R. Fernandez-Lafuente, *Process Biochem.*, 2017, **54**, 81–88.
- 1512 345 S. Peirce, V. Tacias-Pascacio, M. Russo, A. Marzocchella, J. Virgen-Ortíz and R. Fernandez-  
1513 Lafuente, *Molecules*, 2016, **21**, 751.
- 1514 346 L. Fernandez-Lopez, J. J. Virgen-Ortíz, S. G. Pedrero, N. Lopez-Carrobles, B. C. Gorines, C.  
1515 Otero and R. Fernandez-Lafuente, *Biocatal. Biotransformation*, 2017, in press, DOI  
1516 [10.1080/10242422.2016.1278212](https://doi.org/10.1080/10242422.2016.1278212)
- 1517 347 H. Zaak, L. Fernandez-Lopez, C. Otero, M. Sassi and R. Fernandez-Lafuente, in press, DOI  
1518 [10.1016/j.enzmictec.2017.07.001](https://doi.org/10.1016/j.enzmictec.2017.07.001)
- 1519 348 I. Mingarro, C. Abad and L. Braco, *Proc. Natl. Acad. Sci. U. S. A.*, 1995, **92**, 3308–3312.
- 1520 349 F. Peißker and L. Fischer, *Bioorganic Med. Chem.*, 1999, **7**, 2231–2237.
- 1521 350 R. Fernández-Lafuente, C. M. Rosell and J. M. Guisán, *Enzyme Microb. Technol.*, 1998, **23**,  
1522 305–310.
- 1523 351 J. C. S. dos Santos, C. Garcia-Galan, R. C. Rodrigues, H. B. de Sant’Ana, L. R. B. Gonçalves  
1524 and R. Fernandez-Lafuente, *Process Biochem.*, 2014, **49**, 1511–1515.
- 1525 352 R. Torres, C. Ortiz, B. C. C. Pessela, J. M. Palomo, C. Mateo, J. M. Guisán and R. Fernández-  
1526 Lafuente, *Enzyme Microb. Technol.*, 2006, **39**, 167–171.
- 1527 353 F. K. Chu, W. Watorek and F. Maley, *Arch. Biochem. Biophys.*, 1983, **223**, 543–555.
- 1528 354 G. Kern, N. Schülke, F. X. Schmid and R. Jaenicke, *Protein Sci.*, 1992, **1**, 120–131.
- 1529 355 R. Torres, C. Mateo, M. Fuentes, J. M. Palomo, C. Ortiz, R. Fernández-Lafuente, J. M. Guisan,  
1530 A. Tam and M. Daminati, *Biotechnol. Prog.*, 2002, **18**, 1221–1226.
- 1531 356 J. C. S. dos Santos, C. Garcia-Galan, R. C. Rodrigues, H. B. de Sant’Ana, L. R. B. Gonçalves  
1532 and R. Fernandez-Lafuente, *Enzyme Microb. Technol.*, 2014, **60**, 1–8.
- 1533 357 G. Fernández-Lorente, L. Betancor, A. V Carrascosa, J. M. Palomo and J. M. Guisan, *J. Am. Oil*  
1534 *Chem. Soc.*, 2012, **89**, 97–102.

- 1535 358 J. M. Palomo, G. Fernández-Lorente, C. Mateo, M. Fuentes, R. Fernández-Lafuente and J. M.  
1536 Guisan, *Tetrahedron: Asymmetry*, 2002, **13**, 1337–1345.
- 1537 359 G. Fernandez-Lorente, R. Fernández-Lafuente, J. M. Palomo, C. Mateo, A. Bastida, J. Coca, T.  
1538 Haramboure, O. Hernández-Justiz, M. Terreni and J. M. Guisán, in *Journal of Molecular*  
1539 *Catalysis - B Enzymatic*, 2001, vol. 11, pp. 649–656.
- 1540 360 D. S. Rodrigues, A. A. Mendes, M. Filice, R. Fernandez-Lafuente, J. M. Guisan and J. M.  
1541 Palomo, *J. Mol. Catal. B Enzym.*, 2009, **58**, 36–40.
- 1542 361 A. A. Mendes, D. S. Rodrigues, M. Filice, R. Fernandez-Lafuente, J. M. Guisan and J. M.  
1543 Palomo, *Tetrahedron*, 2008, **64**, 10721–10727.
- 1544 362 G. Fernandez-Lorente, M. Filice, M. Terreni, J. M. Guisan, R. Fernandez-Lafuente and J. M.  
1545 Palomo, *J. Mol. Catal. B Enzym.*, 2008, **51**, 110–117.
- 1546 363 J. M. Palomo, M. Filice, R. Fernandez-Lafuente, M. Terreni and J. M. Guisan, *Adv. Synth.*  
1547 *Catal.*, 2007, **349**, 1969–1976.
- 1548 364 J. M. Palomo, R. L. Segura, M. Fuentes, C. C. Ortiz, J. M. Guisán and R. Fernández-Lafuente,  
1549 *Enzyme Microb. Technol.*, 2006, **38**, 429–435.
- 1550 365 J. M. Palomo, G. Fernández-Lorente, C. Mateo, M. Fuentes, J. M. Guisan and R. Fernández-  
1551 Lafuente, *Tetrahedron: Asymmetry*, 2002, **13**, 2653–2659.
- 1552 366 J. M. Palomo, G. Fernandez-Lorente, C. Mateo, C. Ortiz, R. Fernandez-Lafuente and J. M.  
1553 Guisan, *Enzyme Microb. Technol.*, 2002, **31**, 775–783.
- 1554 367 J. Razumiene, A. Vilkanauskite, V. Gureviciene, V. Laurinavicius, N. V. Roznyatovskaya, Y.  
1555 V. Ageeva, M. D. Reshetova and A. D. Ryabov, *J. Organomet. Chem.*, 2003, **668**, 83–90.
- 1556 368 M. K. Ram, M. Adami, S. Paddeu and C. Nicolini, *Nanotechnology*, 2000, **11**, 112–119.
- 1557 369 N. A. Pchelintsev, A. Vakurov and P. A. Millner, *Sensors Actuators, B Chem.*, 2009, **138**, 461–  
1558 466.
- 1559 370 T. Bahar, *Asia-Pacific J. Chem. Eng.*, 2016, **11**, 981–988.
- 1560 371 H. Zheng, J. Zhou, J. Zhang, R. Huang, H. Jia and S. I. Suye, *Microchim. Acta*, 2009, **165**, 109–  
1561 115.
- 1562 372 P. Liu, H. Zheng, P. Nie, Y. Wei, Z. Feng and T. Sun, in *Proceedings of SPIE*, eds. J. Leng, A.  
1563 K. Asundi and W. Ecke, 2009, vol. 7493, p. 74934Y.
- 1564 373 V. Uppada, S. Bhaduri and S. B. Noronha, *Curr. Sci.*, 2014, 106, 946–957.
- 1565 374 X. Ji, Z. Su, P. Wang, G. Ma and S. Zhang, *ACS Nano*, 2015, **9**, 4600–4610.
- 1566 375 S. Velasco-Lozano, A. I. A. I. Benítez-Mateos and F. López-Gallego, *Angew. Chemie Int. Ed.*,  
1567 2017, **56**, 771–775.
- 1568 376 H. Zaak, L. Fernandez-Lopez, C. Otero, M. Sassi, R. Fernandez-Lafuente, *Enzyme Microb.*  
1569 *Technol.*, 2017, **106**, 67–74.

1570 **Figure legends**

1571

1572 **Scheme 1. Standard synthetic route to produce branched-PEI (BPEI) via ring-opening**  
1573 **polymerization of unsubstituted ethylenimine.**

1574 **Scheme 2. Alternative synthetic route to produce branched PEI (BPEI) via ring-opening**  
1575 **isomerization polymerization of substituted 2-oxazolines.**

1576 **Scheme 3. Synthesis of Lineal PEI (LPEI) via ring opening polymerization of 2-ethyl-2-oxazoline**  
1577 **followed by hydrolysis.**

1578 **Figure 1. Reversible immobilization of enzyme via ion exchange.** The support may be reused after  
1579 enzyme inactivation by incubation under more drastic conditions than operation ones.

1580 **Figure 2. Stabilization of multimeric enzymes via multi-subunit immobilization and physical**  
1581 **crosslinking with PEI.** Multimeric enzymes subunits may dissociate during operation (e.g., by  
1582 dilution). Physical crosslinking of the non-immobilized subunits with subunits immobilized on the  
1583 support may prevent this phenomenon.

1584 **Figure 3. Stabilization of multimeric enzymes via multi-subunit crosslinking with PEI and**  
1585 **further immobilization on anion exchangers.** The multimeric enzymes may be coated with PEI to  
1586 prevent subunit dissociation. This composite will be very rich in cationic groups and can be strongly  
1587 immobilized on anionic supports, even if the unmodified enzyme cannot become immobilized on these  
1588 supports.

1589 **Figure 4. Generation of nano-environments around enzyme surface by coating with PEI.** The  
1590 cationic nature of PEI may form a hydrophilic shell surrounding the enzyme molecule, forming  
1591 polymeric salts surrounding the enzyme molecule.



**Figure 5. Change of enzyme properties by physical coating with PEI.** The interaction of an enzyme with a cationic polymer may produce difficulties to the movement of the enzyme or induce conformational changes. These changes should be reversible if the PEI is desorbed.

**Figure 6. Stabilization of enzyme by coating with PEI versus organic solvents:** generation of hydrophilic nano-environments to generate solvent partition.

**Figure 7. Multilayer immobilization using PEI as glue.** After coating the support with a layer of the enzyme, this is activated by coating with PEI and a new layer of enzyme is immobilized. This may be performed continuously until the increment of activity is not compensate by the immobilization of new layers of enzyme (diffusional problems to the inner layers will increase with the production of additional layers of PEI/enzyme).

**Figure 8. Effect of PEI concentration in the final thickness of the polymeric bed.** Using a high concentration of PEI, the support is rapidly coated by many PEI molecules and do not permit to maximize enzyme-support interactions, enabling the formation of thick polymeric beds. Using low concentration of PEI, each PEI molecule may maximize their interactions with the supports because no other PEI molecule will be there avoiding this, forming a thin polymeric bed.

**Figure 9. Immobilization of enzyme in complex reactors using PEI activation of the support walls.** Some bioreactors require a long time to become fixed. In these cases it may be convenient to use reversible strategies of immobilization to avoid the necessity to prepare the reactor anew.

**Figure 10. Reuse of PEI activated supports after enzyme inactivation:** formation of enzyme-support composites with maximized enzyme-support interactions during enzyme inactivation that may complicate to achieve a fully clean support for immobilizing a new batch of protein

**Figure 11. Immobilization of enzymes in PEI coated nano-particles.** The enzyme is immobilized on the surface of the support, that way it is able to interact with large structures. This makes that a

properly oriented enzyme may be active after immobilization even versus solid substrate but also may become inactivated by interaction with hydrophobic surfaces, like gas bubbles.

**Figure 12. Buffering effect of PEI: reduction of the pH gradients.** In some enzyme reactions, e.g hydrolysis of esters at alkaline pH values, a pH gradient may be produced inside the biocatalyst particle if the enzyme activity is high- enough. This pH drop inside the particle in many instances may reduce the enzyme activity or stability. The presence of PEI may be expected to reduce this internal pH gradient due to their high buffering capacity.

**Figure 13. Design of an auto-sufficient glucose biosensor using the multifunctional nature of PEI.** Mediators and GOS are immobilized on the PEI bed to get a glucose biosensor that work without the addition of external mediators.

**Figure 14. PEI coated cells to immobilize enzymes on the cell wall, and their further coimmobilization.** This way, enzymes and cells may act in a synergic way.

**Figure 15. Production of PEI capsules to immobilize enzymes.**

**Figure 16. Use of PEI as enzyme-enzyme glue to co-immobilize two enzymes enabling the reuse of the most stable one.** The first enzyme may be immobilized using optimal protocol to stabilize it, then it is coated with PEI and a second enzyme and less stable may be immobilized on the first activated enzyme. After inactivation of the second enzyme, this may be desorbed and the most stable one reused after immobilizing a new batch of the second enzyme.

**Figure 17. Use of PEI in the preparation of crosslinked enzyme aggregates (CLEAs) of poor Lys enzymes.** Crosslinking may be difficult if the enzyme surface is poor in amino groups. Modification of the enzyme with PEI provide this amino groups and simplify the crosslinking.

**Figure 18. Physical crosslink with PEI of enzyme molecules physically adsorbed on hydrophobic supports.** Lipases immobilized on hydrophobic supports present very good performance but they can be released at high temperature, in presence of organic solvents or detergents, even certain substrates or



product with detergent properties. The coating of the immobilized enzymes with PEI permits the physical intermolecular crosslinking and that way the prevention of enzyme release.

**Figure 19. Stabilization of lipase open form induced by detergents using PEI coating.** Detergents shift the conformational equilibrium of lipases towards the open form, but after washing it, the enzyme recovers the equilibrium. The coating of the lipases in presence of detergent has proved to be useful to keep this detergent induced open form of the lipase.

**Figure 20. Coating of co-immobilized enzymes with PEI permits the coimmobilization of phosphorylated cofactors.**

**Figure 21. Use of PEI to coimmobilize several enzymes and their phosphorylated cofactors.**

1649     **Table 1. Examples of enzyme immobilized on PEI coated supports.**

Enzyme	Support	Application	Reference
Nucleotidase	Polystyrene tube	Determination of hypoxanthine	82
Nuclease P1	weak base anion resin	Hydrolysis of DNA	83
Monoamine oxidase	polypropylene membrane	Oxidation of Tyramine	84
Alkaline phosphatase	Plate-shaped cellulose	Hydrolysis of <i>p</i> -nitrophenyl phosphate	85
Thermophilic $\beta$ -galactosidase	Glyoxyl-agarose beads	Hydrolysis of o-nitrophenyl- $\beta$ -D-galactopyranoside	86
Mesophilic $\beta$ -galactosidase	Glyoxyl-agarose beads	Hydrolysis of o-nitrophenyl- $\beta$ -D-galactopyranoside	87
Mesophilic $\beta$ -galactosidase	Nanofiltration membrane	Production of galactose polymers	88
Mesophilic $\beta$ -galactosidase	Quartz crystal	Hydrolysis of o-nitrophenyl- $\beta$ -D-galactopyranoside	89
Mesophilic $\beta$ -galactosidase	Polymeric membrane	Production of galactose polymers	90
$\beta$ -glucosidase	Cellulose beads	Wine-making and fruit-juice processing	91
$\beta$ -Glucosidase	Glyoxyl-agarose beads	Hydrolysis of cellobiose	92
$\alpha$ -Galactosidase	Glyoxyl-agarose beads	Hydrolysis p-Nitrophenyl- $\alpha$ -D-galactopyranoside	93
$\beta$ -xylosidase	Glyoxyl-agarose beads	Hydrolysis of xylooligosaccharides and production of xylooligosaccharides (X2-X6)	94
Haloalkane dehalogenase	$\gamma$ -alumina beads	Hydrolysis of haloalkane	95
Glucoamylase	Glyoxyl-agarose beads	Hydrolysis of maltose or starch	96
Invertase	Poly(GMA–MMA) beads	Hydrolysis of sucrose	97
Glutaryl Acylase	Epoxy Sepbeads	Hydrolysis of glutaryl-7-ACA	98
Cephalosporin C acylase	Epoxy-LX1000-EPC4	Hydrolysis of cephalosporin C	99
Tyrosinase	Poly(hydroxyethyl methacrylate-co-glycidyl methacrylate) membranes	Transformation of L-Tyr to tyrosinaseo-benzoquinine	100
Tyrosinase	Agar particles, blocks and egg shell	Transformation of L-tyrosine to L-3, 4 dihydroxyphenylalanine	101
Gutamate oxidase	Platinum electrode	Detection of Glu	102

Formate dehydrogenase	Glyoxyl-agarose beads	Oxidation of formic acid, regeneration of NADH	103,104
Penicillin acylase	SiO <sub>2</sub> beads	Hydrolysis of penicillin G	105,106
Alcohol oxidases	Glyoxyl-agarose beads	Ethanol oxidation	107
Glucose oxidase	Screen-printed carbon electrodes	Glucose detection	108
Thermophilic alcohol dehydrogenase	Glyoxyl-agarose beads	Reduction of acetophenone to (S)-(-)-1-phenylethanol	109
Peroxidase	Electrodes with zirconium alcoxide film	Detection of acetaminophen	110
Urease	Egg shell	Detection of urea	111
Laccase	Chitosan	Oxidation of syringaldazine	112
Pectinase	Pulp fiber	Treatment of whitewater from papermaking	113
Cellobiose dehydrogenase	Graphite Electrodes	Cellobiose quantification	114
Pepsin	Epoxy Sepabeads	Hydrolysis of immunoglobulins to produce F(ab') <sub>2</sub> fragments	115
Polygalacturonase	Glyoxyl-agarose beads	Juice clarification	116

1650

1651

1652

1653     **Table 2. Examples of enzyme immobilized on PEI coated supports and treated with**  
1654     **glutaraldehyde.**

Enzyme	Support	Application	Reference
Glucoamylase	Porous glass beads	Hydrolysis of starch	129
Glucose oxidase and catalase	Cotton cloth	Production of spray-dried, sugar-free egg powder	130
Alkaline phosphatase	Silk fabric	Hydrolysis of disodium 4-nitorophenyl phosphate	131
Invertase	Rice husk	Hydrolysis of sucrose	132
Urease	Glass-sealed metal microelectrode	Urea detection	133
β-galactosidase	cotton cloth	Galacto-oligosaccharides production	134, 135
β-galactosidase	Polyether sulfone membrane	Galacto-oligosaccharides production	136
β-galactosidase	Poly (acrylonitrile-	Hydrolysis of lactose	137


	<b>co-methyl methacrylate) poly (AN-co- MMA) Nanofibers</b>		
<b><math>\beta</math>-galactosidase</b>	<b>Flat-sheet ultrafiltration membrane</b>	<b>Galacto-oligosaccharides production</b>	<b>138</b>
<b>D-amino acid oxidase</b>	<b>Glyoxyl agarose beads</b>	<b>Oxidative deamination of Ala and cephalosporin C</b>	<b>139</b>
<b>Lipase</b>	<b>Cotton cloth</b>	<b>Esterification of butanol and butyric acid</b>	<b>140</b>
<b>Lipase</b>	<b>Cotton terry cloth fibrils</b>	<b>Hydrolysis of p-nitrophenyl propionate</b>	<b>141</b>
<b>Lipase</b>	<b>Woolen cloth</b>	<b>Hydrolysis of tributyrin emulsion</b>	<b>142</b>
<b>Lipase</b>	<b>Polyurethane foam</b>	<b>Production of geranyl propionate</b>	<b>143</b>
<b>Urease</b>	<b>Cotton cloth</b>	<b>Hydrolysis of urea</b>	<b>144</b>

1656 **Table 3. Examples of enzyme immobilized on PEI-glutaraldehyde pre-activated supports.**

Enzyme	Support	Application	Reference
trypsin	Silica gel	Hydrolysis of N-cr-benzoyl-L-arginine ethyl ester and N-o-tosyl-L-arginine methyl ester	149
Lactate dehydrogenase	Silica gel	Pyruvate reduction	149
Chymotrypsin	Silica gel	Hydrolysis of N-o-acetyl-L-tyrosine ethyl ester and N-cx-benzoyl-L-tyrosine ethyl ester	149
Glucose oxidase	Nylon	Oxidation of glucose	150
Glucose oxidase	Light-addressable potentiometric sensors	Detection of glucose	151
Chymotrypsin	Light-addressable potentiometric sensors	Detection of peptides	151
Urease	Stain etched porous silico	Detection of urea	152
Urease	Polymethylglutamate membrane	Detection of urea	153
Amyloglucosidase and $\alpha$ -amylase	Different inorganic supports	Hydrolysis of maltose, maltooligosaccharides and soluble starch	154
Amyloglucosidase and $\alpha$ -amylase	Different inorganic supports	Hydrolysis of glycogen	155
Glucose oxidase, malate dehydrogenase	Screen-printed electrodes	Following glucose and malic acid in wine production	156
Amino acid oxidase, protease	Rodinised carbon electrode	Protein detection	157
Glucoamylase	Cotton cloth	Hydrolysis of starch	158
Protease C	Different cloth fibers	Antimicrobial activity	159
Glutaryl acylase	Epoxy SepabedS	Hydrolysis of glutararyl-7-ACA	160
Penicillin G acylase	SiO <sub>2</sub> beads	Hydrolysis of penicillin G	161
Lysozyme and trypsin	Stainless steel	Generation of anti-biofilm surfaces	162
Peroxidase	Electrospun microfibrinous membranes	Degradation of bisphenol A	163
Peroxidase	Aluminum oxide	Oxidation of <i>o</i> -phenylenediamine	164
Trypsin	Aluminum oxide	Hydrolysis of <i>N</i> - $\alpha$ -benzoyl-dl-arginine- <i>p</i> -nitroanilide	164
Invertase	Polyurethane, plast-film	Hydrolysis of sucrose	165


	and ferromagnetic Dacron		
$\beta$ -D-galactosidase	Agar disks	Hydrolysis p-Nitrophenyl-a-D-galactopyranoside	166
Lipase	Polyurethane foam	Production of lauryl laurate	167
Lipase	Woolen fabrics	Stain removal	168
Alcohol oxidase	Electrospun fibers	Determination of ethanol in saliva	169
Dextranucrase	Fe <sup>3+</sup> -cross-linked alginate/carboxymethyl cellulose beads	Dextran synthesis	170
Ppyruvate kinase and L-lactic dehydrogenase	poly(N-isopropylacrylamide) microgel	Transformation of phosphoenol pyruvate on pyruvic acid and ATP and this on lactate with NADH consumption	171


1657




**Figure 1**




**Figure 2**




**Figure 3**

**Figure 4**

**Physically modified enzymes with PEI:**  
The interactions with the polymer may alter enzyme mobility or induce conformational changes altering their catalytic properties




**Figure 5**




Immobilized enzyme

PEI



Organic solvent



Organic solvent







Figure 6




**Figure 7**

**Figure 8**





**Figure 10**




Figure 11







Figure 12



Coimmobilization of GOx and mediators to get a biosensor that not require addition of further compounds to work

Figure 13

**Figure 14**

Coimmobilization of enzymes and cells





Figure 15





**Coimmobilization of two enzymes to reuse the most stable one after inactivation of the less stable enzyme**

**Figure 16**






Figure 17



**Figure 18**



# Active cofactor immobilization on immobilized enzymes: recycling and reuse of cofactors




Figure 20






Figure 21




## Scheme 1





**Scheme 2**

**Scheme 3**

